arXiv:1208.3043v1 [math.PR] 15 Aug 2012

Positive recurrence and transience of multidimensional skip-free reflecting random walks with a background process

Toshihisa Ozawa
Faculty of Business Administration, Komazawa University
1-23-1 Komazawa, Setagaya-ku, Tokyo 154-8525, Japan
E-mail: toshi@komazawa-u.ac.jp

August 16, 2012

Abstract

We consider a discrete-time d-dimensional process $\{\boldsymbol{X}_n\}$ on $\mathbb{Z}_+^d$ with a background process $\{J_n\}$ on a finite set, where the front process $\{\boldsymbol{X}_n\}$ is skip free in all directions. We assume that the joint process $\{\boldsymbol{Y}_n\}=\{(\boldsymbol{X}_n,J_n)\}$ is Markovian and that the transition probabilities of the front process $\{\boldsymbol{X}_n\}$ are modulated depending on the state of the background process $\{J_n\}$. This modulation is space homogeneous in the inside of $\mathbb{Z}_+^d$ and in each boundary face of $\mathbb{Z}_+^d$. We call this model a d-dimensional skip-free reflecting random walk with a background process and obtain sufficient conditions on which it is positive recurrent and on which it is transient.

Keywords: Multidimensional reflecting random walk, positive recurrence, Foster's criterion, quasi-birth-and-death process

Mathematical Subject Classification: 60J10, 60K25

1 Introduction

We consider a discrete-time d-dimensional process $\{\boldsymbol{X}_n\}=\{(X_{1,n},X_{2,n},...,X_{d,n})\}$ on $\mathbb{Z}_+^d$ with a supplemental process $\{J_n\}$ on a finite set, where $\mathbb{Z}_+$ is the set of nonnegative integers. We call the supplemental process a background process. The front process $\{\boldsymbol{X}_n\}$ is skip free in all directions, which means that the increment of each element of the front process takes values in $\{-1,0,1\}$. We assume that the joint process $\{\boldsymbol{Y}_n\}=\{(\boldsymbol{X}_n,J_n)\}$ is Markovian and that the transition probabilities of the front process $\{\boldsymbol{X}_n\}$ are modulated depending on the state of the background process $\{J_n\}$. This modulation is space homogeneous in the inside of $\mathbb{Z}_+^d$ and in each boundary face of $\mathbb{Z}_+^d$. We call this model a d-dimensional skip-free reflecting random walk with a background process (d-dimensional RRW-WBP, in short). Our aim is to exploit sufficient conditions on which the d-dimensional RRW-WBP is positive recurrent and on which it is transient.

The multidimensional RRW-WBP is an extension of the ordinary multidimensional reflecting random walk and it has a lot of potential applications such as those of queueing models; Applications of queueing models spread over various areas, for example, production systems, information network systems and certain service systems, and such a system is often modeled by a multiclass queuing network because it consists of some nodes and contains some kinds of customer. Here we mention one typical example: A single-class d-node generalized Jackson network with Markovian arrivals and phase-type services. The behavior of this model can be represented as a continuous-time

Markov chain $\{\bar{\boldsymbol{Y}}(t)\} = \{(\bar{X}_1(t), \bar{X}_2(t), ..., \bar{X}_d(t), \bar{J}(t))\}$, where $\bar{X}_i(t)$ is the number of customers in the ith node at time t and $\bar{J}(t)$ is the state (phase) of the process combining the Markovian arrival processes and phase-type service processes (see, for example, Neuts [11] and Latouche and Rawaswami [7]). This $\{\bar{\boldsymbol{Y}}(t)\}$ is a continuous-time version of d-dimensional RRW-WBP and, by the uniformization technique, we obtain a discrete-time d-dimensional RRW-WBP, $\{\boldsymbol{Y}_n\}$, that has the same stationary distribution as $\{\bar{\boldsymbol{Y}}(t)\}$ if it exists. This implies that $\{\bar{\boldsymbol{Y}}(t)\}$ is positive recurrent if and only if $\{\boldsymbol{Y}_n\}$ is positive recurrent, and in order to check positive recurrence of the former, it suffices to check that of the latter. Furthermore, in the framework of multidimensional RRW-WBP, we can add some kinds of service discipline and some kinds of interaction between nodes (queues) to the model, by means of Markovian service processes (see, for example, Ozawa [12]). Therefore, understanding positive recurrence and transience of multidimensional RRWs-WBP provides an important basis for analyzing such a queueing model. Of course, stability of queueing networks was studied in a lot of literature in recent decades (see Meyn and Down [9], Dai [3], Bramson [1] and references therein), but in the literature, in order to study the stability, they used stochastic models differed from ours such as fluid limit models.

Recently, asymptotics for the stationary distributions of reflecting random walks were studied in some literature (see, for example, Miyazawa [10] and references therein) and those of two-dimensional quasi-birth-and-death (QBD) processes, which correspond to two-dimensional RRWs-WBP, were studied in Ozawa [13]. Understanding positive recurrence and transience of multidimensional RRWs-WBP provides also a basis for such studies.

Multidimensional RRWs *without a background process* are ordinary multidimensional RRWs, and their positive recurrence and transience were studied by Malyshev [8] and Fayolle et al. [5]. They took the following approach. Let $\mathbb{R}_+$ be the set of nonnegative real numbers and consider a d-dimensional RRW, denoted by $\{\boldsymbol{X}_n\} = \{(X_{1,n}, X_{2,n}, ..., X_{d,n})\}$. First, they defined induced Markov chains, each of which is, roughly speaking, a kind of projection of $\{\boldsymbol{X}_n\}$ onto some boundary face (for a precise definition, see Subsection 4.1) and then defined the second vector field on $\mathbb{R}_+^d$ on which each point of $\mathbb{R}_+^d$ is assigned the mean increment vector of $\{\boldsymbol{X}_n\}$ evaluated by the stationary distribution of a certain induced Markov chain. Considering paths on the second vector field and applying Foster's criterion, they gave sufficient conditions on which the RRW $\{\boldsymbol{X}_n\}$ is positive recurrent and on which it is transient, in terms of certain conditions for the second vector filed. In this paper, we take a similar approach, but *do not use the concept of second vector filed.* Instead of doing so, we consider a certain embedded Markov chain of the d-dimensional RRW-WBP, $\{\boldsymbol{Y}_n\} = \{(\boldsymbol{X}_n, J_n)\}$, and directly apply Foster's criterion to the embedded Markov chain. As a result, we obtain sufficient conditions on which $\{\boldsymbol{Y}_n\}$ is positive recurrent and on which it is transient; each of the sufficient conditions is given by a condition on which there exists at least one fisible solution to a system of linear and quadratic inequalities whose coefficients are given by the elements of the mean increment vectors of the front process $\{\boldsymbol{X}_n\}$ evaluated by the stationary distributions of the induced Markov chains generated from $\{\boldsymbol{Y}_n\}$. Applying these results to two and three-dimensional models, we also obtain the corresponding sufficient conditions in explicit form.

The rest of the paper is organized as follows. In Section 2, the d-dimensional RRW-WBP we consider is described in detail. In Section 3, Foster's criterion and its variations are introduced. In Section 4, sufficient conditions on which the d-dimensional RRW-WBP is positive recurrent and on which it is transient are obtained, and in Section 5, those results are applied to two and three-dimensional models. A simple three-queue model with service interactions is presented in Section 5, as an example of three-dimensional RRW-WBP.

2 Reflecting random walk with a background process

Let d be a positive integer and define a set N as $N = \{1, 2, ..., d\}$. Define a function $\varphi : \mathbb{Z}_+^d \to \mathcal{N}$ as

$$\varphi(\boldsymbol{x}) = N \setminus \{l \in N : x_l = 0\} \text{ for } \boldsymbol{x} = (x_1, x_2, ..., x_d) \in \mathbb{Z}_+^d,$$

where $\mathcal{N}$ is the set of all subsets of N. Note that we treat vectors such as $\boldsymbol{x}$ as column vectors in formulae, except for probability vectors, which we treat as row vectors. For $A \in \mathcal{N}$, let S^A be a finite set, say $S^A = \{1, 2, ..., s^A\}$ for some positive integer s^A. Let a state space $\mathcal{S}$ be defined as

$$\mathcal{S} = \bigcup_{\boldsymbol{x} \in \mathbb{Z}_+^d} \left(\{\boldsymbol{x}\} \times S^{\varphi(\boldsymbol{x})} \right),$$

and consider a time-homogeneous Markov chain $\{\boldsymbol{Y}_n\} = \{(\boldsymbol{X}_n, J_n)\}$ on the state space $\mathcal{S}$; we call the process $\{\boldsymbol{X}_n\} = \{(X_{1,n}, X_{2,n}, \cdots, X_{d,n})\}$ the front process of $\{\boldsymbol{Y}_n\}$ and $\{J_n\}$ the background process of it. From the definition, it can be seen that the transition probabilities of the front process are modulated depending on the state of the background process. We call the process $\{\boldsymbol{Y}_n\}$ a d-dimensional reflecting random walk with a background process (d-dimensional RRW-WBP, in short) and denote it by $\mathcal{L}$. Throughout the paper, we assume the following condition.

Assumption 2.1. *$\mathcal{L}$ is irreducible.*

Furthermore, we assume that individual processes $\{X_{l,n}\}$, $l \in N$, are skip free and introduce a kind of space-homogeneous structure of transition probabilities, as follows. For $A \in \mathcal{N}$, let $\mathcal{B}^A$ be the boundary face defined as

$$\mathcal{B}^A = \{(\boldsymbol{x}, i) \in \mathcal{S} : \varphi(\boldsymbol{x}) = A\};$$

$\mathcal{B}^N$ is the interior space of $\mathcal{S}$, but we also call it a boundary face. Transition probabilities are space-homogeneous in each boundary face and they are represented as, for $(\boldsymbol{x}, i) \in \mathcal{S}$, $\boldsymbol{z} \in \{-1, 0, 1\}^d$ such that $\boldsymbol{x} + \boldsymbol{z} \in \mathbb{Z}_+^d$ and $j \in S^{\varphi(\boldsymbol{x}+\boldsymbol{z})}$,

$$P(\boldsymbol{Y}_{n+1} = (\boldsymbol{x} + \boldsymbol{z}, j) \,|\, \boldsymbol{Y}_{n+1} = (\boldsymbol{x}, i)) = p_{\boldsymbol{z}}^{\varphi(\boldsymbol{x})\, \varphi(\boldsymbol{x}+\boldsymbol{z})}(i, j). \tag{2.1}$$

Let $\boldsymbol{\xi}_n = (\xi_{1,n}, \xi_{2,n}, ..., \xi_{d,n})$ be the vector of increments defined as $\boldsymbol{\xi}_n = \boldsymbol{X}_n - \boldsymbol{X}_{n-1}$. By the assumption of skip-free property, we have $\boldsymbol{\xi}_n \in \{-1, 0, 1\}^d$. For $\boldsymbol{y} \in \mathcal{S}$, we denote by $\boldsymbol{\alpha}(\boldsymbol{y})$ the vector of the conditional mean increments given that the chain is in state $\boldsymbol{y}$, i.e.,

$$\boldsymbol{\alpha}(\boldsymbol{y}) = (\alpha_1(\boldsymbol{y}), \alpha_2(\boldsymbol{y}), \cdots, \alpha_d(\boldsymbol{y})),\ \alpha_l(\boldsymbol{y}) = E(\xi_{l,n+1} \,|\, \boldsymbol{Y}_n = \boldsymbol{y}),\ l \in N.$$

3 Preliminaries

We present some existing results which are used for proving theorems after. Note that, in this section, notations are defined independently of other sections.

Let $\{Z_n\}$ be a discrete-time Markov chain on a countable set $\mathcal{S}$ and assume that it is time-homogenous and irreducible. The following proposition is called Foster's criterion and it gives a necessary and sufficient condition on which a Markov chain is positive recurrent.

Proposition 3.1 (Foster's criterion)**.** *The Markov chain* $\{Z_n\}$ *is positive recurrent if and only if there exist a positive number* ε*, a finite subset* $\mathcal{V} \subset \mathcal{S}$ *and a lower bounded real function* $f(i)$*,* $i \in \mathcal{S}$*, such that*

$$E(f(Z_{n+1}) - f(Z_n) \,|\, Z_n = i) \le -\varepsilon, \;\; i \in \mathcal{S} \setminus \mathcal{V}, \tag{3.1}$$

$$E(f(Z_{n+1}) \,|\, Z_n = i) < \infty, \;\; i \in \mathcal{V}. \tag{3.2}$$

Exclusively divide the state space $\mathcal{S}$ into a finite number of nonempty subsets, say $\mathcal{V}_k$, $k = 1, 2, ..., m_S$, and define a function $\tau(i)$ as

$$\tau(i) = T_k \text{ if } i \in \mathcal{V}_k,$$

where T_k, $k \in \{1, 2, ..., m_S\}$, is a positive integer. Furthermore, define a strictly increasing time sequence $\{t_n\}$ as

$$t_{n+1} = t_n + \tau(Z_{t_n}), \; n \ge 0, \;\; t_0 = 0,$$

and consider a stochastic process $\{\tilde{Z}_n\}$ defined by $\tilde{Z}_n = Z_{t_n}$, $n \ge 0$. Obviously this process $\{\tilde{Z}_n\}$ is a kind of embedded Markov chain of $\{Z_n\}$. If the embedded Markov chain is irreducible, then a recurrence time for each state in the embedded Markov chain is always greater than or equal to the corresponding recurrence time for the same state in the original Markov chain. This implies that if $\{\tilde{Z}_n\}$ is positive recurrent then $\{Z_n\}$ is also positive recurrent, and hence we obtain the following corollary, which is a simple version of Theorem 2.2.4 of Fayolle et al. [5]; also see Theorem 3.2 of Fayolle [4] and Proposition 4.5 of Bramson [1].

Corollary 3.1. *Assume that the embedded Markov chain* $\{\tilde{Z}_n\}$ *is irreducible. If there exist a positive number* ε*, a finite subset* $\mathcal{V} \subset \mathcal{S}$ *and a lower bounded real function* $f(i)$*,* $i \in \mathcal{S}$*, such that*

$$E(f(\tilde{Z}_{n+1}) - f(\tilde{Z}_n) \,|\, \tilde{Z}_n = i) \le -\varepsilon, \;\; i \in \mathcal{S} \setminus \mathcal{V}, \tag{3.3}$$

$$E(f(\tilde{Z}_{n+1}) \,|\, \tilde{Z}_n = i) < \infty, \;\; i \in \mathcal{V}, \tag{3.4}$$

then the Markov chain $\{Z_n\}$ *is positive recurrent.*

In order to prove that a Markov chain is transient, we can use the following proposition, which is a simplified version of Theorem 2.2.7 of Fayolle et al. [5].

Proposition 3.2. *Assume that the embedded Markov chain* $\{\tilde{Z}_n\}$ *is irreducible. If there exist a real function* $f(i)$*,* $i \in \mathcal{S}$*, and positive numbers* ε*,* c *and* b *such that, for* $\mathcal{A} = \{i \in \mathcal{S} : f(i) > c\}$*,*

(i) $\mathcal{A} \ne \emptyset$*,* $\mathcal{A}^C \ne \emptyset$*,*

(ii) $E(f(\tilde{Z}_{n+1}) - f(\tilde{Z}_n) \,|\, \tilde{Z}_n = i) \ge \varepsilon, \;\; i \in \mathcal{A}$*, and*

(iii) the inequality $|f(j) - f(i)| > b$ *implies* $P(Z_1 = j \,|\, Z_0 = i) = 0$*,*

then the Markov chain $\{Z_n\}$ *is transient.*

4 Positive recurrence and transience

In this section, we deal with a d-dimensional RRW with a background process, $\mathcal{L} = \{\boldsymbol{Y}_n\} = \{(\boldsymbol{X}_n, J_n)\}$, introduced in Section 2, and derive sufficient conditions on which $\mathcal{L}$ is positive recurrent and on which it is transient. For the purpose, we introduce three kinds of Markov chain generated from $\mathcal{L}$, including induced Markov chains, and then obtain approximations for the expected time-averaged increments of the front process $\{\boldsymbol{X}_n\}$. This approximation plays an essential role in deriving main results.

4.1 Markov chains generated from $\mathcal{L}$

Hereafter, we use the following notations. For $A \subset N$, we denote by $\boldsymbol{x}^A$ a vector in $\mathbb{Z}_+^{|A|}$ whose elements are indexed in A, i.e., $\boldsymbol{x}^A = (x_l,\, l \in A)$, and by $\boldsymbol{2}^A$ the $|A|$-dimensional vector of 2's whose elements are indexed in A, where $|A|$ is the cardinality of set A. We also denote by $(\boldsymbol{x}^A, \boldsymbol{y}^{N\setminus A})$ the combination of $\boldsymbol{x}^A$ and $\boldsymbol{y}^{N\setminus A}$; the l^{th} element of $(\boldsymbol{x}^A, \boldsymbol{y}^{N\setminus A})$ is x_l if $l \in A$ and y_l if $l \in N \setminus A$. For example, when $d = 5$ and $A = \{1, 3\}$, we have $\boldsymbol{x}^A = (x_1, x_3)$, $\boldsymbol{y}^{N\setminus A} = (y_2, y_4, y_5)$ and $(\boldsymbol{x}^A, \boldsymbol{y}^{N\setminus A}) = (x_1, y_2, x_3, y_4, y_5)$. We also use this notation for expressing a part of a given vector; For $\boldsymbol{x} = (x_l, l \in N)$ and $A \subset N$, we denote $(x_l, l \in A)$ by $\boldsymbol{x}^A$ and represent $\boldsymbol{x}$ as $\boldsymbol{x} = (\boldsymbol{x}^A, \boldsymbol{x}^{N\setminus A}) = (\boldsymbol{x}^{N\setminus A}, \boldsymbol{x}^A)$.

4.1.1 Expanded Markov chain

We consider a Markov chain obtained from $\mathcal{L}$ by removing some boundaries. For $A \subset N$ such that $A \neq \emptyset$, let $\bar{\mathcal{S}}_A$ be a state space defined as

$$\bar{\mathcal{S}}_A = \bigcup_{\boldsymbol{x}^{N\setminus A} \in \mathbb{Z}_+^{d-|A|}} \; \bigcup_{\boldsymbol{x}^A \in \mathbb{Z}^{|A|}} \left(\{(\boldsymbol{x}^{N\setminus A}, \boldsymbol{x}^A)\} \times S^{\varphi(\boldsymbol{x}^{N\setminus A}, \boldsymbol{2}^A)} \right), \tag{4.1}$$

and consider a skip-free partially-reflecting random walk with a background process on $\bar{\mathcal{S}}_A$, denoted by $\{(\bar{\boldsymbol{X}}_n, \bar{J}_n)\} = \{((\bar{\boldsymbol{X}}_n^{N\setminus A}, \bar{\boldsymbol{X}}_n^A), \bar{J}_n)\}$, whose transition probabilities are given by, for $(\boldsymbol{x}, i) = ((\boldsymbol{x}^{N\setminus A}, \boldsymbol{x}^A), i) \in \bar{\mathcal{S}}_A$ and $\boldsymbol{z} = (\boldsymbol{z}^{N\setminus A}, \boldsymbol{z}^A) \in \{-1, 0, 1\}^d$ such that $(\boldsymbol{x} + \boldsymbol{z}, j) \in \bar{\mathcal{S}}_A$,

$$P\big((\bar{\boldsymbol{X}}_{n+1}, \bar{J}_{n+1}) = (\boldsymbol{x} + \boldsymbol{z}, j) \,|\, (\bar{\boldsymbol{X}}_n, \bar{J}_n) = (\boldsymbol{x}, i)\big) = p_{\boldsymbol{z}}^{\varphi(\boldsymbol{x}^{N\setminus A}, \boldsymbol{2}^A)\, \varphi(\boldsymbol{x}^{N\setminus A} + \boldsymbol{z}^{N\setminus A}, \boldsymbol{2}^A)}(i, j), \tag{4.2}$$

where $p_{\boldsymbol{z}}^{\varphi(\boldsymbol{x})\, \varphi(\boldsymbol{x}+\boldsymbol{z})}(i, j)$ is given in expression (2.1). We call $\{(\bar{\boldsymbol{X}}, \bar{J}_n)\}$ an expanded Markov chain of $\mathcal{L}$ concerning A and denote it by $\bar{\mathcal{L}}_A$.

4.1.2 Induced Markov chain

For $A \subset N$ such that $A \neq \emptyset$, we define a state space $\mathcal{S}^A$ as

$$\mathcal{S}^A = \bigcup_{\boldsymbol{x}^{N\setminus A} \in \mathbb{Z}_+^{d-|A|}} \left(\{\boldsymbol{x}^{N\setminus A}\} \times S^{\varphi(\boldsymbol{x}^{N\setminus A}, \boldsymbol{2}^A)} \right), \tag{4.3}$$

and introduce a Markov chain $\mathcal{L}^A$ on $\mathcal{S}^A$, called an induced Markov chain, according to Malyshev [8] and Fayolle et al. [5]; $\mathcal{L}^A$ is the $(d - |A|)$-dimensional RRW with a background process, $\{(\hat{\boldsymbol{X}}_n^{N\setminus A}, \hat{J}_n)\}$, whose transition probabilities are given by, for $(\boldsymbol{x}^{N\setminus A}, i) \in \mathcal{S}^A$ and $\boldsymbol{z}^{N\setminus A} \in$

$\{-1,0,1\}^{d-|A|}$ such that $\boldsymbol{x}^{N\setminus A}+\boldsymbol{z}^{N\setminus A}\in\mathbb{Z}_+^{d-|A|}$ and for $j\in S^{\varphi(\boldsymbol{x}^{N\setminus A}+\boldsymbol{z}^{N\setminus A},\boldsymbol{2}^A)}$,

$$\begin{aligned} &P\Big((\hat{\boldsymbol{X}}_{n+1}^{N\setminus A},\hat{J}_{n+1})=(\boldsymbol{x}^{N\setminus A}+\boldsymbol{z}^{N\setminus A},j)\,\big|\,(\hat{\boldsymbol{X}}_{n}^{N\setminus A},\hat{J}_{n})=(\boldsymbol{x}^{N\setminus A},i)\Big) \\ &\qquad=\sum_{\boldsymbol{z}^A\in\{-1,0,1\}^{|A|}} p_{(\boldsymbol{z}^{N\setminus A},\boldsymbol{z}^A)}^{\varphi(\boldsymbol{x}^{N\setminus A},\boldsymbol{2}^A)\,\varphi(\boldsymbol{x}^{N\setminus A}+\boldsymbol{z}^{N\setminus A},\boldsymbol{2}^A)}(i,j). \end{aligned}\tag{4.4}$$

Throughout the paper, we assume the following condition.

Assumption 4.1. *For all $A\subset N$ such that $A\neq\emptyset$, $\mathcal{L}^A$ is irreducible.*

For $A\subset N$ such that $A\neq\emptyset$, if $\mathcal{L}^A$ is positive recurrent, then we denote by $\boldsymbol{\pi}^A=(\pi^A(\boldsymbol{x}^{N\setminus A},i),\,(\boldsymbol{x}^{N\setminus A},i)\in\mathcal{S}^A)$ its stationary distribution. Note that $\mathcal{L}^N$ is a finite Markov chain and it is always positive recurrent. For $\mathcal{L}^A$ being positive recurrent, we define the d-dimensional mean increment vector, $\boldsymbol{a}(A)=(a_l(A),\,l\in N)$, by

$$a_l(A)=\sum_{(\boldsymbol{x}^{N\setminus A},i)\in\mathcal{S}^A}\alpha_l((\boldsymbol{x}^{N\setminus A},\boldsymbol{2}^A),i)\,\pi^A(\boldsymbol{x}^{N\setminus A},i),\ \ l\in N.\tag{4.5}$$

From the definition, it can be seen that $a_l(A)=0$ for all $l\in N\setminus A$.

Remark 4.1. *From the definition of induced Markov chain and that of extended Markov chain, it can be seen that, for each $A\subset N$, $A\neq\emptyset$, the probability law of the induced Markov chain $\mathcal{L}^A$ is identical to that of the partial process (marginal process) $\{(\bar{\boldsymbol{X}}_n^{N\setminus A},\bar{J}_n)\}$ of the extended Markov chain $\bar{\mathcal{L}}_A$. In other wards, $\mathcal{L}^A$ is the projection of $\bar{\mathcal{L}}_A$ onto the boundary face $\mathcal{B}^{N\setminus A}$.*

4.1.3 Embedded Markov chain

For $A\subset N$ such that $A\neq\emptyset$, let K_A be a positive integer and assume that, for $A,B\subset N$, if $|A|>|B|$ then $K_A<K_B$. We inductively define, for $A\subset N$, a subset $\mathcal{V}_A\subset\mathcal{S}$ as follows (for the case of $d=2$, see Fig. 1).

Step 1 Set $k:=d$ and $\mathcal{V}_\emptyset:=\mathcal{S}$.

Step 2 For $A\subset N$ such that $|A|=k$, set $\mathcal{V}_A:=\mathcal{V}_\emptyset\cap\{(\boldsymbol{x},i)\in\mathcal{S}:x_l\geq K_A,\ l\in A\}$. Set $\mathcal{V}_\emptyset:=\mathcal{V}_\emptyset\setminus\Big(\bigcup_{A\subset N,\,|A|=k}\mathcal{V}_A\Big)$ and $k:=k-1$.

Step 3 If $k>0$ then go to Step 2, otherwise we obtain $\mathcal{V}_A$ for all $A\subset N$.

It is obvious that $\bigcup_{A\subset N}\mathcal{V}_A=\mathcal{S}$ and that $\mathcal{V}_A\cap\mathcal{V}_B=\emptyset$ for all $A,B\subset N$ such that $A\neq B$. Thus, $\{\mathcal{V}_A:A\subset N\}$ is a partition of $\mathcal{S}$. We also see that $\mathcal{V}_\emptyset$ is finite. For $\boldsymbol{y}\in\mathcal{S}$, define a function $\tau(\boldsymbol{y})$ as

$$\tau(\boldsymbol{y})=T_A \text{ if } \boldsymbol{y}\in\mathcal{V}_A \text{ for some } A\subset N,$$

where $T_A,\ A\subset N$, are positive integers. We assume that $T_\emptyset=1$ and that, for $A\subset N$ such that $A\neq\emptyset$, $T_A<K_A$. Furthermore, define a time sequence $\{t_n\}$ as $t_{n+1}=t_n+\tau(\boldsymbol{Y}_{t_n})$, $t_0=0$, and a Markov chain $\{\tilde{\boldsymbol{Y}}_n\}$ as $\tilde{\boldsymbol{Y}}_n=\boldsymbol{Y}_{t_n}$, $n\geq 0$. We call $\{\tilde{\boldsymbol{Y}}_n\}$ the embedded Markov chain of $\mathcal{L}$ and denote it by $\tilde{\mathcal{L}}$. Since the original Markov chain $\mathcal{L}$ is irreducible, we can make the embedded Markov chain $\tilde{\mathcal{L}}$ irreducible by appropriately setting parameters T_A, $A\subset N$, $A\neq\emptyset$, and this

does not restrict the values of the parameters from above. Hence, hereafter we assume that the parameters T_A, $A \subset N\, A \neq \emptyset$, satisfy that condition. By Corollary 3.1, if there exists a set of parameters $\{(K_A, T_A) : A \subset N, A \neq \emptyset\}$ such that the condition of the corollary holds, then the Markov chain $\mathcal{L}$ is positive recurrent.

Remark 4.2. *Since the front process of the RRW-WBP $\{\boldsymbol{Y}_n\}$ is skip free in each direction, it can be seen that, for $A \subset N$ such that $A \neq \emptyset$, if $\tilde{\boldsymbol{Y}}_k \in \mathcal{V}_A$ then the Markov chain $\{\boldsymbol{Y}_n\}$ does not visit any state on the boundary face $\mathcal{B}^{N\setminus A}$ during the time interval $[t_k, t_{k+1}]$. Thus, in stochastic sense, the process $\{\boldsymbol{Y}_n\}$ behaves just like the expanded Markov chain $\bar{\mathcal{L}}_A$ during that time interval.*



Figure 1: Partition of the state space ($d = 2$).

4.2 Time average of increments

Here we consider the expected time-averaged increments of the front process $\{\boldsymbol{X}_n\}$ and obtain their approximations, which will be used as an important tool for deriving our main results.

Consider a d-dimensional RRW-WBP $\{\boldsymbol{Y}_n\} = \{(\boldsymbol{X}_n, J_n)\}$ on state space $\mathcal{S}$. Let $\mathcal{N}_p$ be the index set of the positive-recurrent induced Markov chains, i.e.,

$$\mathcal{N}_p = \{A \subset N : A \neq \emptyset,\ \mathcal{L}^A \text{ is positive recurrent}\}.$$

Since $\mathcal{L}^N$ is always positive recurrent, $\mathcal{N}_p$ is not empty. For $A \subset N$ such that $A \neq \emptyset$, let $\mathcal{N}^A$ be the set of the subsets of N including all elements of A, i.e., $\mathcal{N}^A = \{B \subset N : A \subset B\}$, and $\mathcal{N}_p^A$ be defined as $\mathcal{N}_p^A = \mathcal{N}^A \cap \mathcal{N}_p$. Note that if $\mathcal{L}^A$ is positive recurrent, then $A \in \mathcal{N}_p^A$. For $\boldsymbol{y} \in \mathcal{S}$, define the vector of the expected time-averaged increments, $\boldsymbol{g}(\boldsymbol{y})$, as

$$\boldsymbol{g}(\boldsymbol{y}) = (g_l(\boldsymbol{y}),\, l \in N), \quad g_l(\boldsymbol{y}) = E\biggl(\frac{1}{\tau(\boldsymbol{y})} \sum_{n=1}^{\tau(\boldsymbol{y})} \xi_{l,n} \,\bigg|\, \boldsymbol{Y}_0 = \boldsymbol{y}\biggr),\ l \in N,$$

where $\xi_{l,n} = X_{l,n} - X_{l,n-1}$. The following proposition asserts that $\boldsymbol{g}(\boldsymbol{y})$ can be approximated by a linear combination of $\boldsymbol{\alpha}(A)$, $A \in \mathcal{N}_p$, the mean increment vectors evaluated by the stationary distributions of the positive-recurrent induced Markov chains.

Proposition 4.1. *For any $\varepsilon > 0$, there exist the set of parameters $\{(T_A, K_A) : A \subset N,\ A \neq \emptyset\}$ satisfying the conditions mentioned in Subsection 4.1.3 and substochastic vectors $(p_A(B),\ B \in \mathcal{N}_p^A)$, $A \subset N$, $A \neq \emptyset$, such that, for every $\boldsymbol{y} \notin \mathcal{V}_\emptyset$, if $\boldsymbol{y} \in \mathcal{V}_A$ for some $A \subset N$, then we have*

$$\Bigl| g_l(\boldsymbol{y}) - \sum_{B \in \mathcal{N}_p^A} p_A(B)\, a_l(B) \Bigr| < \varepsilon,\ l \in N, \tag{4.6}$$

where some of $p_A(B)$, $B \in \mathcal{N}_p^A$, may be zero and K_A can be taken sufficiently largely, independently of T_A. Especially, if $\mathcal{L}^A$ is positive recurrent, we have

$$\bigl| g_l(\boldsymbol{y}) - a_l(A) \bigr| < \varepsilon,\ l \in N. \tag{4.7}$$

For simplicity, we denote $g_l(\boldsymbol{y})$ having property (4.6) by

$$g_l(\boldsymbol{y}) \approx \sum_{B \in \mathcal{N}_p^A} p_A(B)\, a_l(B).$$

Proof. For $A \subset N$ such that $A \neq \emptyset$, let $\{\bar{\boldsymbol{Y}}_n\} = \{(\bar{\boldsymbol{X}}_n, \bar{J}_n)\}$ be the extended Markov chain $\bar{\mathcal{L}}_A$ and define the conditional expected time-average of increments as, for $l \in N$ and $\boldsymbol{y} \in \bar{\mathcal{S}}_A$,

$$\bar{g}_l(A, \boldsymbol{y}) = E\biggl(\frac{1}{T_A} \sum_{n=1}^{T_A} \bar{\xi}_{l,n} \,\Big|\, (\bar{\boldsymbol{X}}_0, \bar{J}_0) = \boldsymbol{y} \biggr),$$

where $\bar{\xi}_{l,n} = \bar{X}_{l,n} - \bar{X}_{l,n-1}$. By Remark 4.2, if $\boldsymbol{y} \in \mathcal{V}_A$, then we have $g_l(\boldsymbol{y}) = \bar{g}_l(A, \boldsymbol{y})$. Therefore, we evaluate $\bar{g}_l(A, \boldsymbol{y})$ instead of $g_l(\boldsymbol{y})$. For $\boldsymbol{y}_0 \in \mathcal{V}_A$,

$$\begin{aligned}
\bar{g}_l(A, \boldsymbol{y}_0) &= \frac{1}{T_A} \sum_{n=0}^{T_A-1} \sum_{\boldsymbol{y}_n \in \bar{\mathcal{S}}_A} E\bigl(\bar{\xi}_{l,n+1} \,|\, (\bar{\boldsymbol{X}}_n, \bar{J}_n) = \boldsymbol{y}_n\bigr) P\bigl((\bar{\boldsymbol{X}}_n, \bar{J}_n) = \boldsymbol{y}_n \,|\, (\bar{\boldsymbol{X}}_0, \bar{J}_0) = \boldsymbol{y}_0\bigr) \\
&= \frac{1}{T_A} \sum_{n=0}^{T_A-1} \sum_{(\boldsymbol{x}_n^{N\setminus A}, j_n) \in \mathcal{S}^A} \sum_{\boldsymbol{x}_n^A \in \mathbb{Z}^{|A|}} E\Bigl(\bar{\xi}_{l,n+1} \,|\, ((\bar{\boldsymbol{X}}_n^{N\setminus A}, \bar{\boldsymbol{X}}_n^A), \bar{J}_n) = ((\boldsymbol{x}_n^{N\setminus A}, \boldsymbol{2}^A), j_n)\Bigr) \\
&\qquad \times P\Bigl(((\bar{\boldsymbol{X}}_n^{N\setminus A}, \bar{\boldsymbol{X}}_n^A), \bar{J}_n) = ((\boldsymbol{x}_n^{N\setminus A}, \boldsymbol{x}_n^A), j_n) \,|\, (\bar{\boldsymbol{X}}_0, \bar{J}_0) = \boldsymbol{y}_0\Bigr) \\
&= \frac{1}{T_A} \sum_{n=0}^{T_A-1} \sum_{(\boldsymbol{x}_n^{N\setminus A}, j_n) \in \mathcal{S}^A} \alpha_l\bigl((\boldsymbol{x}_n^{N\setminus A}, \boldsymbol{2}^A), j_n\bigr) \\
&\qquad \times P\Bigl((\bar{\boldsymbol{X}}_n^{N\setminus A}, \bar{J}_n) = (\boldsymbol{x}_n^{N\setminus A}, j_n) \,|\, (\bar{\boldsymbol{X}}_0, \bar{J}_0) = \boldsymbol{y}_0\Bigr) \\
&= E\biggl(\frac{1}{T_A} \sum_{n=0}^{T_A-1} \alpha_l\bigl((\bar{\boldsymbol{X}}_n^{N\setminus A}, \boldsymbol{2}^A), \bar{J}_n\bigr) \,\Big|\, (\bar{\boldsymbol{X}}_0, \bar{J}_0) = \boldsymbol{y}_0 \biggr)
\end{aligned} \tag{4.8}$$

where $\boldsymbol{y}_n$ is expressed as $\boldsymbol{y}_n = (\boldsymbol{x}_n, j_n) = ((\boldsymbol{x}_n^{N\setminus A}, \boldsymbol{x}_n^A), j_n)$. By Remark 4.1, the probability law of the partial process $\{(\bar{\boldsymbol{X}}_n^{N\setminus A}, \bar{J}_n)\}$ is identical to that of the induced Markov chain $\mathcal{L}^A$. Thus, by the ergodic theorem for Markov chains, if $\mathcal{L}^A$ is positive recurrent, then for a sufficiently large T_A, we obtain

$$\bigl| \bar{g}_l(A, \boldsymbol{y}_0) - a_l(A) \bigr| < \varepsilon,\ \ l \in N,$$

where $a_l(A)$ is given by expression (4.5); this is identical to inequality (4.6) and K_A can be taken arbitrarily largely.

Next, we prove the proposition in the case where a concerning induced Markov chain is not positive recurrent, by induction. Consider the induced Markov chain $\mathcal{L}^N$, where $|N| = d$. Since $\mathcal{L}^N$ is positive recurrent, inequality (4.6) holds for some $T_N > 0$ and K_N can be taken arbitrarily largely. Suppose that there exists some parameter set $\{(T_A, K_A) : A \subset N,\ |A| \ge k\}$ such that, for all $A \subset N$, $|A| \ge k$, inequality (4.6) holds. Let A be a subset of N satisfying $|A| = k-1$. If the induced Markov chain $\mathcal{L}^A$ is positive recurrent, then inequality (4.6) holds for some $T_A > 0$ and K_A can be taken arbitrarily largely. If it is not, we introduce an embedded Markov chain generated from the expanded Markov chain $\bar{\mathcal{L}}_A$, denoted by $\{\tilde{\boldsymbol{Y}}^\dagger_n\} = \{(\tilde{\boldsymbol{X}}^\dagger_n, \tilde{J}^\dagger_n)\}$. For $B \in \mathcal{N}^A$, let $\bar{\mathcal{V}}_B$ be a subset of $\bar{\mathcal{S}}_A$, obtained by expanding $\mathcal{V}_B$ as

$$\bar{\mathcal{V}}_B = \{((\boldsymbol{x}^{N\setminus A}, \boldsymbol{x}^A), i) \in \bar{\mathcal{S}}_A : ((\boldsymbol{x}^{N\setminus A}, \boldsymbol{y}^A), i) \in \mathcal{V}_B \text{ for some } \boldsymbol{y}^A \in \mathbb{Z}_+^{|A|}\}.$$

From the definition, it can be seen that $\{\bar{\mathcal{V}}_B,\ B \in \mathcal{N}^A\}$ is a partition of $\bar{\mathcal{S}}_A$. For $\boldsymbol{y} \in \bar{\mathcal{S}}_A$, define a function $\bar{\tau}(\boldsymbol{y})$ as

$$\bar{\tau}(\boldsymbol{y}) = \begin{cases} T_B & \text{if } \boldsymbol{y} \in \bar{\mathcal{V}}_B \text{ for some } B \in \mathcal{N}^A \text{ such that } B \ne A, \\ 1 & \text{otherwise,} \end{cases}$$

and a time sequence $\{\bar{t}_n\}$ as $\bar{t}_0 = 0$, $\bar{t}_{n+1} = \bar{t}_n + \bar{\tau}(\bar{\boldsymbol{Y}}_{t_n})$, $n \ge 0$. The embedded Markov chain $\{\tilde{\boldsymbol{Y}}^\dagger_n\}$ is defined as $\tilde{\boldsymbol{Y}}^\dagger_n = \bar{\boldsymbol{Y}}_{\bar{t}_n}$, $n \ge 0$. By the assumption of induction, we have, for $B \in \mathcal{N}^A$ such that $B \ne A$ and for $\boldsymbol{y} \in \bar{\mathcal{V}}_B$,

$$\frac{1}{T_B} E\big(\tilde{X}^\dagger_{l,1} - \tilde{X}^\dagger_{l,0} \,|\, \tilde{\boldsymbol{Y}}^\dagger_0 = \boldsymbol{y}\big) = \bar{g}_l(B, \boldsymbol{y}) \approx \sum_{C \in \mathcal{N}^B_p} p_B(C)\, a_l(C). \tag{4.9}$$

For $\boldsymbol{y} \in \bar{\mathcal{V}}_A$, $\bar{g}_l(A, \boldsymbol{y})$ is represented as

$$\begin{aligned} \bar{g}_l(A, \boldsymbol{y}) &= \frac{1}{T_A} E\big(\bar{X}_{l,T_A} - \bar{X}_{l,0} \,|\, \bar{\boldsymbol{Y}}_0 = \boldsymbol{y}\big) \\ &= \frac{1}{T_A} \sum_{n=1}^{T_A} E\Big(1(\bar{t}_n \le T_A)\, (\tilde{X}^\dagger_{l,n} - \tilde{X}^\dagger_{l,n-1}) \\ &\qquad + 1(\bar{t}_{n-1} \le T_A < \bar{t}_n)\, (\bar{X}_{l,T_A} - \tilde{X}^\dagger_{l,n-1}) \,\big|\, \tilde{\boldsymbol{Y}}^\dagger_0 = \boldsymbol{y}\Big) \\ &= \sum_{B \in \mathcal{N}^A} \phi_1(B) + \phi_2, \end{aligned} \tag{4.10}$$

where $1(\cdot)$ is an indicator function and, for $B \in \mathcal{N}^A$,

$$\phi_1(B) = \frac{1}{T_A} \sum_{n=1}^{T_A} \sum_{\boldsymbol{y}' \in \bar{\mathcal{V}}_B} E\Big(1(\bar{t}_n \le T_A)\, (\tilde{X}^\dagger_{l,n} - \tilde{X}^\dagger_{l,n-1}) \,\big|\, \tilde{\boldsymbol{Y}}^\dagger_{n-1} = \boldsymbol{y}'\Big)\, P\big(\tilde{\boldsymbol{Y}}^\dagger_{n-1} = \boldsymbol{y}' \,|\, \tilde{\boldsymbol{Y}}^\dagger_0 = \boldsymbol{y}\big)$$

and

$$\phi_2 = \frac{1}{T_A} \sum_{n=1}^{T_A} E\Big(1(\bar{t}_{n-1} \le T_A < \bar{t}_n)\, (\bar{X}_{l,T_A} - \tilde{X}^\dagger_{l,n-1}) \,\big|\, \tilde{\boldsymbol{Y}}^\dagger_0 = \boldsymbol{y}\Big).$$

Because of the skip-free property of the chain and the assumption of $\bar{\tau}(\boldsymbol{y}) = 1$ for $\boldsymbol{y} \in \bar{\mathcal{V}}_A$, we obtain

$$\begin{aligned}
\phi_1(A) &\le \frac{1}{T_A} \sum_{n=1}^{T_A} \sum_{\boldsymbol{y}' \in \bar{\mathcal{V}}_A} E\Big(\bar{X}_{l,n} - \bar{X}_{l,n-1} \,\big|\, \bar{\boldsymbol{Y}}_{n-1} = \boldsymbol{y}' \Big)\, P\big(\bar{\boldsymbol{Y}}_{n-1} = \boldsymbol{y}' \,|\, \bar{\boldsymbol{Y}}_0 = \boldsymbol{y} \big) \\
&\le \frac{1}{T_A} \sum_{n=1}^{T_A} P\big(\bar{\boldsymbol{Y}}_{n-1} \in \bar{\mathcal{V}}_A \,|\, \bar{\boldsymbol{Y}}_0 = \boldsymbol{y} \big), \qquad (4.11)
\end{aligned}$$

and this implies that $\phi_1(A) \approx 0$ for a sufficiently large T_A because the induced Markov chain $\mathcal{L}^A$ is not positive recurrent and the set $\{(\boldsymbol{x}^{N\setminus A}, i) \in \mathcal{S}^A : ((\boldsymbol{x}^{N\setminus A}, \boldsymbol{y}^A), i) \in \bar{\mathcal{V}}_A \text{ for some } \boldsymbol{y} \in \mathbb{Z}^{|A|}\}$ is finite. We also have

$$\phi_2 \le \frac{\max_{B \in \mathcal{N}^A, B \ne A} T_B}{T_A}, \qquad (4.12)$$

and this implies that $\phi_2 \approx 0$ for a sufficiently large T_A. Since $\bar{t}_n = \bar{t}_{n-1} + \bar{\tau}(\tilde{\boldsymbol{Y}}^\dagger_{n-1})$, we have, for $B \in \mathcal{N}^A$, $B \ne A$,

$$\begin{aligned}
\phi_1(B) &= \frac{1}{T_A} \sum_{n=1}^{T_A} \sum_{\boldsymbol{y}' \in \bar{\mathcal{V}}_B} E\Big(\tilde{X}^\dagger_{l,n} - \tilde{X}^\dagger_{l,n-1} \,\big|\, \tilde{\boldsymbol{Y}}^\dagger_{n-1} = \boldsymbol{y}',\, \bar{t}_{n-1} \le T_A - \bar{\tau}(\boldsymbol{y}') \Big) \\
&\qquad\qquad \times P\big(\tilde{\boldsymbol{Y}}^\dagger_{n-1} = \boldsymbol{y}',\, \bar{t}_{n-1} + \bar{\tau}(\tilde{\boldsymbol{Y}}^\dagger_{n-1}) \le T_A \,|\, \tilde{\boldsymbol{Y}}^\dagger_0 = \boldsymbol{y} \big) \\
&= \frac{1}{T_A} \sum_{n=1}^{T_A} \sum_{\boldsymbol{y}' \in \bar{\mathcal{V}}_B} T_B\, \bar{g}_l(B, \boldsymbol{y}')\, P\big(\tilde{\boldsymbol{Y}}^\dagger_{n-1} = \boldsymbol{y}',\, \bar{t}_n \le T_A \,|\, \tilde{\boldsymbol{Y}}^\dagger_0 = \boldsymbol{y} \big),
\end{aligned}$$

and this implies that

$$\Bigg| \phi_1(B) - q_A(B) \sum_{C \in \mathcal{N}^B_p} p_B(C)\, a_l(C) \Bigg| < q_A(B)\, \varepsilon, \qquad (4.13)$$

where

$$q_A(B) = \frac{T_B}{T_A} \sum_{n=1}^{T_A} P\big(\tilde{\boldsymbol{Y}}^\dagger_{n-1} \in \bar{\mathcal{V}}_B,\, \bar{t}_n \le T_A \,|\, \tilde{\boldsymbol{Y}}^\dagger_0 = \boldsymbol{y} \big).$$

Since $\bar{\tau}(\boldsymbol{y}') = 1$ for $\boldsymbol{y}' \in \bar{\mathcal{V}}_A$, we define $q_A(A)$ as

$$q_A(A) = \frac{1}{T_A} \sum_{n=1}^{T_A} P\big(\tilde{\boldsymbol{Y}}^\dagger_{n-1} \in \bar{\mathcal{V}}_A,\, \bar{t}_n \le T_A \,|\, \tilde{\boldsymbol{Y}}^\dagger_0 = \boldsymbol{y} \big).$$

Then, we have

$$\begin{aligned}
\sum_{B \in \mathcal{N}^A} q_A(B) \quad &= \frac{1}{T_A} \sum_{n=1}^{T_A} \sum_{B \in \mathcal{N}^A} E\bigg(\bar{\tau}(\tilde{\boldsymbol{Y}}^\dagger_{n-1})\, 1(\tilde{\boldsymbol{Y}}^\dagger_{n-1} \in \bar{\mathcal{V}}_B)\, 1(\bar{t}_n \le T_A) \,\bigg|\, \tilde{\boldsymbol{Y}}^\dagger_0 = \boldsymbol{y} \bigg) \\
&= \frac{1}{T_A} E\bigg(\sum_{n=1}^{T_A} \bar{\tau}(\tilde{\boldsymbol{Y}}^\dagger_{n-1})\, 1(\bar{t}_n \le T_A) \,\bigg|\, \tilde{\boldsymbol{Y}}^\dagger_0 = \boldsymbol{y} \bigg),
\end{aligned}$$

and this implies

$$1-\max_{B\in\mathcal{N}^A,B\neq A}\frac{T_B}{T_A}\le\sum_{B\in\mathcal{N}^A}q_A(B)\le 1. \tag{4.14}$$

On the other hand, $q_A(A)$ satisfies

$$q_A(A)\le\frac{1}{T_A}\sum_{n=1}^{T_A}P\big(\bar{\boldsymbol{Y}}_{n-1}\in\bar{\mathcal{V}}_A\,|\,\bar{\boldsymbol{Y}}_0=\boldsymbol{y}\big), \tag{4.15}$$

and this implies that $q_A(A)\approx 0$ for a sufficiently large T_A because of the same reason as that why $\phi_1(A)\approx 0$ for a sufficiently large T_A. Therefore, we obtain, for a sufficiently large T_A,

$$0<1-\max_{B\in\mathcal{N}^A,B\neq A}\frac{T_B}{T_A}-q_A(A)\le\sum_{B\in\mathcal{N}^A,B\neq A}q_A(B)\le 1, \tag{4.16}$$

and from this, we see that vector $(q_A(B),\,B\in\mathcal{N}^A,B\neq A)$ is substochastic. For $C\in\mathcal{N}_p^A$, let $p_A(C)$ be defined as

$$p_A(C)=\sum_{B\in\mathcal{N}^A,B\neq A}q_A(B)\,p_B(C),$$

then vector $(p_A(C),\,C\in\mathcal{N}_p^A)$ is also substochastic. As a result, from expression (4.10), we obtain, for a sufficiently large T_A,

$$\begin{aligned}&\Bigg|\bar{g}_l(A,\boldsymbol{y})-\sum_{C\in\mathcal{N}_p^A}p_A(C)\,a_l(C)\Bigg|\\&\qquad\le\sum_{B\in\mathcal{N}^A,B\neq A}\Bigg|\phi_1(B)-q_A(B)\sum_{C\in\mathcal{N}_p^B}p_B(C)\,a_l(C)\Bigg|+|\phi_1(A)|+|\phi_2|<\varepsilon,\end{aligned} \tag{4.17}$$

where K_A can be taken arbitrarily largely; This completes the proof. □

4.3 Main results

Let $U=(\boldsymbol{u}_j)=(u_{i,j})$ be a $d\times d$ symmetric matrix, where $\boldsymbol{u}_j$ is the jth column composing U, and define a set of $d\times d$ positive definite matrices, $\mathcal{U}$, as

$$\mathcal{U}=\Big\{U=(\boldsymbol{u}_j):U\text{ is positive definite and }\langle\boldsymbol{a}(A),\boldsymbol{u}_j\rangle<0\text{ for all }A\in\mathcal{N}_p\text{ and all }j\in A\Big\},$$

where $\langle\boldsymbol{x}_1,\boldsymbol{x}_2\rangle$ is the inner product of vectors $\boldsymbol{x}_1,\boldsymbol{x}_2\in\mathbb{R}^d$ and $\boldsymbol{a}(A)$ is the mean increment vector evaluated by the stationary distribution of the induced Markov chain $\mathcal{L}^A$ which is positive recurrent. The following theorem provides a sufficient condition on which the d-dimensional RRW-WBP is positive recurrent.

Theorem 4.1. *The d-dimensional RRW with a background process is positive recurrent if* $\mathcal{U}\neq\emptyset$.

Proof. Let $U = (\boldsymbol{u}_j) = (u_{i,j})$ be a $d \times d$ positive definite matrix and introduce the following quadratic function as a test function:

$$f(\boldsymbol{y}) = f(\boldsymbol{x}, i) = \frac{1}{2} \langle \boldsymbol{x}, U\boldsymbol{x} \rangle, \ \boldsymbol{y} = (\boldsymbol{x}, i) \in \mathcal{S}. \tag{4.18}$$

In order to prove the theorem, it suffices to show that there exists a positive definite matrix U satisfying the condition of Corollary 3.1. Let $\{\tilde{\boldsymbol{Y}}_n\}$ be the embedded Markov chain of $\{\boldsymbol{Y}_n\}$ defined in Subsection 4.1.3. We have $\tilde{\boldsymbol{X}}_{n+1} = \tilde{\boldsymbol{X}}_n + \sum_{k=1}^{\tau(\boldsymbol{Y}_{t_n})} \boldsymbol{\xi}_{t_n+k}$, where $\boldsymbol{\xi}_k = \boldsymbol{X}_k - \boldsymbol{X}_{k-1}$, and the difference between $f(\tilde{\boldsymbol{Y}}_{n+1})$ and $f(\tilde{\boldsymbol{Y}}_n)$ satisfies

$$\begin{aligned} f(\tilde{\boldsymbol{Y}}_{n+1}) - f(\tilde{\boldsymbol{Y}}_n) &= \frac{1}{2} \left\langle \sum_{k=1}^{\tau(\boldsymbol{Y}_{t_n})} \boldsymbol{\xi}_{t_n+k}, U \sum_{k=1}^{\tau(\boldsymbol{Y}_{t_n})} \boldsymbol{\xi}_{t_n+k} \right\rangle + \left\langle \sum_{k=1}^{\tau(\boldsymbol{Y}_{t_n})} \boldsymbol{\xi}_{t_n+k}, U\tilde{\boldsymbol{X}}_n \right\rangle \\ &\le c_0(\tau(\boldsymbol{Y}_{t_n}), U) + \left\langle \sum_{k=1}^{\tau(\boldsymbol{Y}_{t_n})} \boldsymbol{\xi}_{t_n+k}, U\tilde{\boldsymbol{X}}_n \right\rangle, \end{aligned} \tag{4.19}$$

where

$$c_0(\tau(\boldsymbol{Y}_{t_n}), U) = \frac{1}{2} \tau(\boldsymbol{Y}_{t_n})^2 \max_{\boldsymbol{x},\boldsymbol{y} \in \{-1,0,1\}^d} \langle \boldsymbol{x}, U\boldsymbol{y} \rangle < \infty$$

and we use the assumption that the RRW-WBP is skip free in all directions. Hence we obtain, for $\boldsymbol{y} = (\boldsymbol{x}, i) \in \mathcal{S}$,

$$\begin{aligned} E(f(\tilde{\boldsymbol{Y}}_{n+1}) - f(\tilde{\boldsymbol{Y}}_n) \,|\, \tilde{\boldsymbol{Y}}_n = \boldsymbol{y}) &\le c_0(\tau(\boldsymbol{y}), U) + E\biggl(\Bigl\langle \sum_{k=1}^{\tau(\boldsymbol{y})} \boldsymbol{\xi}_{t_n+k}, U\boldsymbol{x} \Bigr\rangle \,\Big|\, \tilde{\boldsymbol{Y}}_n = \boldsymbol{y} \biggr) \\ &= c_0(\tau(\boldsymbol{y}), U) + \tau(\boldsymbol{y}) \langle \boldsymbol{g}(\boldsymbol{y}), U\boldsymbol{x} \rangle. \end{aligned} \tag{4.20}$$

Let A be a nonempty subset of N. For $\boldsymbol{y} = (\boldsymbol{x}, i) \in \mathcal{V}_A$, we have $x_l \le \max_{B \in \mathcal{N}^A, B \ne A} K_B$ for $l \in N \setminus A$, and thus

$$\begin{aligned} &E(f(\tilde{\boldsymbol{Y}}_{n+1}) - f(\tilde{\boldsymbol{Y}}_n) \,|\, \tilde{\boldsymbol{Y}}_n = \boldsymbol{y}) \\ &\quad \le c_0(T_A, U) + T_A \sum_{l \in N \setminus A} \bigl| \langle \boldsymbol{g}(\boldsymbol{y}), \boldsymbol{u}_l \rangle \bigr| \max_{B \in \mathcal{N}^A, B \ne A} K_B + T_A \sum_{l \in A} \langle \boldsymbol{g}(\boldsymbol{y}), \boldsymbol{u}_l \rangle \, x_l, \end{aligned} \tag{4.21}$$

where we use the fact that $\langle \boldsymbol{g}(\boldsymbol{y}), U\boldsymbol{x} \rangle = \sum_{l=1}^{d} \langle \boldsymbol{g}(\boldsymbol{y}), \boldsymbol{u}_l \rangle \, x_l$. The first and second terms on the right hand side of inequality (4.21) are bounded and do not depend on the value of K_A. On the other hand, we have $x_l \ge K_A$ for all $l \in A$. Thus, if $\langle \boldsymbol{g}(\boldsymbol{y}), \boldsymbol{u}_l \rangle < 0$ for all $l \in A$, there exists a sufficiently large K_A such that $E(f(\tilde{\boldsymbol{Y}}_{n+1}) - f(\tilde{\boldsymbol{Y}}_n) \,|\, \tilde{\boldsymbol{Y}}_n = \boldsymbol{y}) \le -\varepsilon$ for some $\varepsilon > 0$. Therefore, by Corollary 3.1, in order to prove the theorem, it suffice to show that there exists a set of parameters $\{(T_A, K_A) : A \subset N,\ N \ne \emptyset\}$ such that, for all $A \subset N$, $A \ne \emptyset$, and for all $l \in A$ and $\boldsymbol{y} \in \mathcal{V}_A$, we have $\langle \boldsymbol{g}(\boldsymbol{y}), \boldsymbol{u}_l \rangle < 0$; it is obvious that $\mathcal{V}_\emptyset$ is finite and $E(f(\tilde{\boldsymbol{Y}}_{n+1}) \,|\, \tilde{\boldsymbol{Y}}_n = \boldsymbol{y}) < \infty$ for all $\boldsymbol{y} \in \mathcal{V}_\emptyset$.

Let U satisfy the condition of the theorem. That is, for all $A \in \mathcal{N}_p$ and for all $l \in A$, we have $\langle \boldsymbol{a}(A), \boldsymbol{u}_l \rangle < 0$. By Proposition 4.1, there exists a set of parameters $\{(T_A, K_A) : A \subset N,\ A \ne \emptyset\}$

such that, for $A \subset N$, $A \neq \emptyset$, and for $\boldsymbol{y} \in \mathcal{V}_A$ and $l \in A$,

$$
\begin{aligned}
\langle \boldsymbol{g}(\boldsymbol{y}), \boldsymbol{u}_l \rangle &\le \sum_{B \in \mathcal{N}_p^A} p_A(B) \, \langle \boldsymbol{a}(B), \boldsymbol{u}_l \rangle + \sum_{i \in N} \left| g_i(\boldsymbol{y}) - \sum_{B \in \mathcal{N}_p^A} p_A(B) \, a_i(B) \right| |u_{il}| \\
&\le \sum_{B \in \mathcal{N}_p^A} p_A(B) \, \langle \boldsymbol{a}(B), \boldsymbol{u}_l \rangle + \varepsilon' \sum_{i \in N} |u_{il}|, \qquad (4.22)
\end{aligned}
$$

where $\varepsilon' > 0$ can be taken arbitrarily small. Thus, we obtain $\langle \boldsymbol{g}(\boldsymbol{y}), \boldsymbol{u}_l \rangle < 0$. Since K_A can be taken arbitrarily largely in Proposition 4.1, this completes the proof. $\square$

Next we consider a condition for the RRW-WBP to be transient. For $A \subset N$ such that $A \neq \emptyset$, let $\bar{\mathcal{N}}^A$ and $\bar{\mathcal{N}}_p^A$ be index sets of induced Markov chains, defined as

$$
\bar{\mathcal{N}}^A = \bigcup_{l \in A} \mathcal{N}^{\{l\}}, \quad \bar{\mathcal{N}}_p^A = \bar{\mathcal{N}}^A \cap \mathcal{N}_p,
$$

and let $\mathcal{W}_A$ be a set of d-dimensional vectors, defined as

$$
\begin{aligned}
\mathcal{W}_A = \Big\{ \boldsymbol{w} = (\boldsymbol{w}^A, \boldsymbol{w}^{N \setminus A}) \in \mathbb{R}^d : \boldsymbol{w}^A > \boldsymbol{0}^A, \ \boldsymbol{w}^{N \setminus A} < \boldsymbol{0}^{N \setminus A}, \\
\text{and } \langle \boldsymbol{a}(B), \boldsymbol{w} \rangle > 0 \text{ for all } B \in \bar{\mathcal{N}}_p^A \Big\},
\end{aligned}
$$

where $\boldsymbol{0} = (\boldsymbol{0}^A, \boldsymbol{0}^{N \setminus A})$ is a vector of 0's with a suitable dimension. A sufficient condition on which the d-dimensional RRW-WBP is transient is given by the following theorem.

Theorem 4.2. *The d-dimensional RRW with a background process, $\{\boldsymbol{Y}_n\} = \{(\boldsymbol{X}_n, J_n)\}$, is transient if $\mathcal{W}_A \neq \emptyset$ for some $A \subset N$, $A \neq \emptyset$.*

Proof. Suppose that there exists $A \subset N$, $A \neq \emptyset$, satisfying $\mathcal{W}_A \neq \emptyset$ and let $\boldsymbol{w} = (\boldsymbol{w}^A, \boldsymbol{w}^{N \setminus A})$ be an element of $\mathcal{W}_A$. Define the following linear function as a test function:

$$
f(\boldsymbol{y}) = f(\boldsymbol{x}, i) = \langle \boldsymbol{x}, \boldsymbol{w} \rangle, \ \boldsymbol{y} = (\boldsymbol{x}, i) \in \mathcal{S}.
$$

Since the front process of $\{\boldsymbol{Y}_n\}$ is skip-free in all directions, we have

$$
|f(\boldsymbol{Y}_{n+1}) - f(\boldsymbol{Y}_n)| = \left| \langle \boldsymbol{X}_{n+1} - \boldsymbol{X}_n, \boldsymbol{w} \rangle \right| \le \sum_{l \in N} |w_l|, \qquad (4.23)
$$

and this implies that if $|f(\boldsymbol{y}') - f(\boldsymbol{y})| > \sum_{l \in N} |w_l|$ then $P(\boldsymbol{Y}_1 = \boldsymbol{y}' \,|\, \boldsymbol{Y}_0 = \boldsymbol{y}) = 0$. Thus, we have $b = \sum_{l \in N} |w_l|$ satisfying condition (iii) of Proposition 3.2. For a positive integer c, define a subset of $\mathcal{S}$, $\mathcal{A}$, as

$$
\mathcal{A} = \{ \boldsymbol{y} = (\boldsymbol{x}, i) \in \mathcal{S} : f(\boldsymbol{y}) = \langle \boldsymbol{x}, \boldsymbol{w} \rangle > c \}.
$$

Since $\boldsymbol{w}^A > \boldsymbol{0}^A$ and $\boldsymbol{w}^{N \setminus A} < \boldsymbol{0}^{N \setminus A}$, we have, for $\boldsymbol{y} = (\boldsymbol{x}, i) \in \mathcal{A}$, $\langle \boldsymbol{x}^A, \boldsymbol{w}^A \rangle > c$. Thus, we can set c so large that $\mathcal{V}_B \cap \mathcal{A} = \emptyset$ for all $B \subset N \setminus A$, and this implies that

$$
\mathcal{A} \subset \mathcal{S} \setminus \bigcup_{B \subset N \setminus A} \mathcal{V}_B = \bigcup_{B \in \bar{\mathcal{N}}^A} \mathcal{V}_B.
$$

Hence, for all $\boldsymbol{y} \in \mathcal{A}$, there exists $B \in \bar{\mathcal{N}}^A$ such that $\boldsymbol{y} \in \mathcal{V}_B$ and, by Proposition 4.1, we obtain, for the $\boldsymbol{y}$,

$$\begin{aligned} E\Big(f(\tilde{\boldsymbol{Y}}_{n+1}) - f(\tilde{\boldsymbol{Y}}_n) \,\big|\, \tilde{\boldsymbol{Y}}_n = \boldsymbol{y}\Big) &= E\Big(\langle \tilde{\boldsymbol{X}}_{n+1} - \tilde{\boldsymbol{X}}_n, \boldsymbol{w}\rangle \,\big|\, \tilde{\boldsymbol{Y}}_n = \boldsymbol{y}\Big) \\ &= \tau(\boldsymbol{y})\, \langle \boldsymbol{g}(\boldsymbol{y}), \boldsymbol{w}\rangle \\ &\approx T_B \sum_{C \in \mathcal{N}_p^B} p_B(C)\, \langle \boldsymbol{a}(C), \boldsymbol{w}\rangle > 0, \end{aligned}$$

where we use the fact that $\mathcal{N}_p^B \subset \bar{\mathcal{N}}_p^A$ because of $B \cap A \neq \emptyset$. This leads to condition (ii) of Proposition 3.2 and complets the proof. $\square$

Remark 4.3. *In the proof of the theorem, $\mathcal{A}$ is a half space of $\mathcal{S}$ separated by the hyperplane $\langle \boldsymbol{x}, \boldsymbol{w}\rangle = c$, where $\boldsymbol{w}$ is a normal vector of the hyperplane. Hence, the condition of the theorem means that the expected time-averaged increment vector of any point $\boldsymbol{y} \in \mathcal{A}$, $\boldsymbol{g}(\boldsymbol{y})$, does not cross the hyperplane. This well works for two-dimensional RRWs-WBP, but does not for a certain class of three-dimensional RRW-WBP, as mentioned in the next section.*

5 Low dimensional cases

In this section, applying Theorems 4.1 and 4.2, we obtain specific conditions on which the d-dimensional RRW with a background process is positive recurrent and on which it is transient, when $d = 2$ and 3. Since a one-dimensional RRW with a background process is an ordinary discrete-time quasi-birth-and-death process (QBD process), a necessary and sufficient condition on which the QBD process is positive recurrent is well known (see, for example, Neuts [11] and Latouche and Rawaswami [7]); We can derive the same result by Theorems 4.1 and 4.2, but it is straightforward and we omit it in the paper.

5.1 Two dimensional case

Consider the case of $d = 2$. Let $\{\boldsymbol{Y}_n\} = \{(\boldsymbol{X}_n, J_n)\}$ be a two-dimensional RRW with a background process, which is denoted by $\mathcal{L}$. We assume that $\mathcal{L}$ is irreducible and that all the induced Markov chains generated from $\mathcal{L}$ are also irreducible. We have $N = \{1, 2\}$ and, for $\boldsymbol{x} = (x_1, x_2) \in \mathbb{Z}_+^2$,

$$\varphi(\boldsymbol{x}) = \begin{cases} \emptyset & \text{if } x_1 = 0 \text{ and } x_2 = 0, \\ \{1\} & \text{if } x_1 > 0 \text{ and } x_2 = 0, \\ \{2\} & \text{if } x_1 = 0 \text{ and } x_2 > 0, \\ N & \text{otherwise.} \end{cases}$$

The state space $\mathcal{S}$ is given by

$$\mathcal{S} = \Big(\{0\} \times \{0\} \times \mathcal{S}^{\emptyset}\Big) \cup \Big(\{0\} \times \mathbb{N} \times \mathcal{S}^{\{1\}}\Big) \cup \Big(\mathbb{N} \times \{0\} \times \mathcal{S}^{\{2\}}\Big) \cup \big(\mathbb{N} \times \mathbb{N} \times \mathcal{S}^{N}\big).$$

For $A, B \subset N$ and $z_1, z_2 \in \{-1, 0, 1\}$, we define a matrix of transition probabilities, $P_{z_1,z_2}^{A,B}$, as

$$P_{z_1,z_2}^{A,B} = \Big(p_{(z_1,z_2)}^{A,B}(i,j),\ i \in S^A,\ j \in S^B\Big),$$

where $p^{A,B}_{(z_1,z_2)}(i,j)$ is given in expression (2.1). We assume $P^{A,B}_{z_1,z_2}$ to be a zero matrix for impossible transitions; for example, we have $P^{N,\{1\}}_{1,1}=O$. We abbreviate $P^{A,A}_{z_1,z_2}$ to $P^{A}_{z_1,z_2}$ and denote the sum of them with respect to z_1 and/or z_2 by

$$P^{A,B}_{*,z_2}=\sum_{z_1\in\{-1,0,1\}}P^{A,B}_{z_1,z_2},\quad P^{A,B}_{z_1,*}=\sum_{z_2\in\{-1,0,1\}}P^{A,B}_{z_1,z_2},\quad P^{A,B}_{*,*}=\sum_{z_1,z_2\in\{-1,0,1\}}P^{A,B}_{z_1,z_2}.$$

The induced Markov chain $\mathcal{L}^N$ is an irreducible finite Markov chain, whose transition probability matrix is given by $P^N_{*,*}$. Hence $\mathcal{L}^N$ is positive recurrent and we denote by $\boldsymbol{\pi}^N$ its stationary distribution. The mean increment vector $\boldsymbol{a}(N)=(a_i(N),i=1,2)$ is given as

$$a_1(N)=\boldsymbol{\pi}^N_{*,*}\big(-P^N_{-1,*}+P^N_{1,*}\big)\,\mathbf{1},\quad a_2(N)=\boldsymbol{\pi}^N_{*,*}\big(-P^N_{*,-1}+P^N_{*,1}\big)\,\mathbf{1},$$

where $\mathbf{1}$ is a column vector of 1's with a suitable dimension. The induced Markov chains $\mathcal{L}^{\{1\}}$ and $\mathcal{L}^{\{2\}}$ are irreducible QBD processes whose transition probability matrices, denoted by $P^{\{1\}}$ and $P^{\{2\}}$, respectively, are given in block form as follows:

$$P^{\{1\}}=\begin{pmatrix} P^{\{1\}}_{*,0} & P^{\{1\},N}_{*,1} & & \\ P^{N,\{1\}}_{*,-1} & P^N_{*,0} & P^N_{*,1} & \\ & P^N_{*,-1} & P^N_{*,0} & P^N_{*,1} \\ & \ddots & \ddots & \ddots \end{pmatrix},\quad P^{\{2\}}=\begin{pmatrix} P^{\{2\}}_{0,*} & P^{\{2\},N}_{1,*} & & \\ P^{N,\{2\}}_{-1,*} & P^N_{0,*} & P^N_{1,*} & \\ & P^N_{-1,*} & P^N_{0,*} & P^N_{1,*} \\ & \ddots & \ddots & \ddots \end{pmatrix}.$$

By Theorem 7.2.3 of Latouche and Rawaswami [7], for $l\in\{1,2\}$, $\mathcal{L}^{\{l\}}$ is positive recurrent if and only if $a_{3-l}(N)<0$; By Theorem 6.2.1 of Latouche and Rawaswami [7], if $\mathcal{L}^{\{l\}}$ is positive recurrent, its stationary distribution $\boldsymbol{\pi}^{\{l\}}$ is given as, in matrix geometric form,

$$\boldsymbol{\pi}^{\{l\}}=(\boldsymbol{\pi}^{\{l\}}_k,k\in\mathbb{Z}_+),\quad \boldsymbol{\pi}^{\{l\}}_k=\boldsymbol{\pi}^{\{l\}}_k(R^{\{l\}})^{k-1},\ k\ge 2,$$

where $R^{\{l\}}$ is the rate matrix of $\mathcal{L}^{\{l\}}$. Hence, the mean increment vectors $\boldsymbol{a}(\{1\})=(a_i(\{1\}),i=1,2)$ and $\boldsymbol{a}(\{2\})=(a_i(\{2\}),i=1,2)$ are given by

$$\begin{aligned}
a_1(\{1\})&=\boldsymbol{\pi}^{\{1\}}_0\big(-P^{\{1\}}_{-1,0}-P^{\{1\},N}_{-1,1}+P^{\{1\}}_{1,0}+P^{\{1\},N}_{1,1}\big)\,\mathbf{1}\\
&\quad+\boldsymbol{\pi}^{\{1\}}_1\big(-P^{N,\{1\}}_{-1,-1}-P^N_{-1,0}-P^N_{-1,1}+P^{N,\{1\}}_{1,-1}+P^N_{1,0}+P^N_{1,1}\big)\,\mathbf{1}\\
&\quad+\boldsymbol{\pi}^{\{1\}}_2\big(I-R^{\{1\}}\big)^{-1}\big(-P^N_{-1,*}+P^N_{1,*}\big)\,\mathbf{1},\\
a_2(\{1\})&=0,\quad a_1(\{2\})=0,\\
a_2(\{2\})&=\boldsymbol{\pi}^{\{2\}}_0\big(-P^{\{2\}}_{0,-1}-P^{\{2\},N}_{1,-1}+P^{\{2\}}_{0,1}+P^{\{2\},N}_{1,1}\big)\,\mathbf{1}\\
&\quad+\boldsymbol{\pi}^{\{2\}}_1\big(-P^{N,\{2\}}_{-1,-1}-P^N_{0,-1}-P^N_{1,-1}+P^{N,\{2\}}_{-1,1}+P^N_{0,1}+P^N_{1,1}\big)\,\mathbf{1}\\
&\quad+\boldsymbol{\pi}^{\{2\}}_2\big(I-R^{\{2\}}\big)^{-1}\big(-P^N_{*,-1}+P^N_{*,1}\big)\,\mathbf{1}.
\end{aligned}$$

Classification of two-dimensional RRWs with a back ground process is given by the following theorem.

Theorem 5.1. *(C1) When $a_1(N)<0$ and $a_2(N)<0$, the 2D-RRW-WBP $\{\mathbf{Y}_n\}$ is*

(a) positive recurrent if $a_1(\{1\})<0$ and $a_2(\{2\})<0$;

(b) transient if either $a_1(\{1\}) > 0$ *or* $a_2(\{2\}) > 0$.

(C2) When $a_1(N) > 0$ *and* $a_2(N) < 0$*, the 2D-RRW-WBP* $\{\boldsymbol{Y}_n\}$ *is*

(a) positive recurrent if $a_1(\{1\}) < 0$*;*

(b) transient if $a_1(\{1\}) > 0$.

(C3) When $a_1(N) < 0$ *and* $a_2(N) > 0$*, the 2D-RRW-WBP* $\{\boldsymbol{Y}_n\}$ *is*

(a) positive recurrent if $a_2(\{2\}) < 0$*;*

(b) transient if $a_2(\{2\}) > 0$.

(C4) When $a_1(N) > 0$ *and* $a_2(N) > 0$*, the 2D-RRW-WBP* $\{\boldsymbol{Y}_n\}$ *is transient.*

Proof. In (C1), we have $\mathcal{N}_p = \{\{1\}, \{2\}, N\}$. First we prove (C1)-(a). In order for a square matrix $U = (\boldsymbol{u}_1, \boldsymbol{u}_2) = (u_{i,j}, i, j = 1, 2)$ to be positive definite, it suffices that $u_{11} > 0$ and $\det U > 0$, where we assume $u_{12} = u_{21}$. Hence, if we find a solution to the following system of inequalities, then the solution is an element of $\mathcal{U}$ in Theorem 4.1:

$$\begin{aligned}
&\langle \boldsymbol{a}(\{1\}), \boldsymbol{u}_1 \rangle = a_1(\{1\})\, u_{11} < 0, \quad \langle \boldsymbol{a}(\{2\}), \boldsymbol{u}_2 \rangle = a_2(\{2\})\, u_{22} < 0, \\
&\langle \boldsymbol{a}(N), \boldsymbol{u}_1 \rangle = a_1(N)\, u_{11} + a_2(N)\, u_{12} < 0, \quad \langle \boldsymbol{a}(N), \boldsymbol{u}_2 \rangle = a_1(N)\, u_{12} + a_2(N)\, u_{22} < 0, \\
&u_{11} > 0, \quad u_{11}u_{22} - u_{12}^2 > 0.
\end{aligned}$$

It is an easy task; For example, that $u_{11} = u_{22} = 1$ and $u_{12} = 1/2$ is such a solution. Thus, $\mathcal{U}$ is not empty and, by Theorem 4.1, the 2D-RRW-WBP is positive recurrent. Next we prove (C1)-(b). Consider the case of $a_1(\{1\}) > 0$. We have $\bar{\mathcal{N}}^{\{1\}} = \{\{1\}, N\}$ and $\bar{\mathcal{N}}_p^{\{1\}} = \{\{1\}, N\}$. Hence, if we find a solution $\boldsymbol{w} = (w_1, w_2)$ to the following system of inequalities, then the solution is an element of $\mathcal{W}_{\{1\}}$ in Theorem 4.2:

$$\langle \boldsymbol{a}(\{1\}), \boldsymbol{w} \rangle = a_1(\{1\})\, w_1 > 0, \quad \langle \boldsymbol{a}(N), \boldsymbol{w} \rangle = a_1(N)\, w_1 + a_2(N)\, w_2 > 0, \quad w_1 > 0,\ w_2 < 0.$$

It is such a solution that $w_1 = 1$ and $w_2 = -a_1(N)/a_2(N) - \delta < 0$ for some $\delta > 0$. Thus, $\mathcal{W}_{\{1\}}$ is not empty and, by Theorem 4.2, the 2D-RRW-WBP is transient. The same result is analogously obtained in the case of $a_2(\{2\}) > 0$.

In (C2), we have $\mathcal{N}_p = \{\{1\}, N\}$. First we prove (C2)-(a). If we find a solution to the following system of inequalities, then the solution is an element of $\mathcal{U}$ in Theorem 4.1:

$$\begin{aligned}
&\langle \boldsymbol{a}(\{1\}), \boldsymbol{u}_1 \rangle = a_1(\{1\})\, u_{11} < 0, \quad \langle \boldsymbol{a}(N), \boldsymbol{u}_1 \rangle = a_1(N)\, u_{11} + a_2(N)\, u_{12} < 0, \\
&\langle \boldsymbol{a}(N), \boldsymbol{u}_2 \rangle = a_1(N)\, u_{12} + a_2(N)\, u_{22} < 0, \quad u_{11} > 0, \quad u_{11}u_{22} - u_{12}^2 > 0.
\end{aligned}$$

This is equivalent to

$$0 < -\frac{a_1(N)}{a_2(N)}\, u_{11} < u_{12} < -\frac{a_2(N)}{a_1(N)}\, u_{22}, \quad u_{11} > 0, \quad u_{22} > 0, \quad u_{11}u_{22} > u_{12}^2,$$

and taking u_{22} to be sufficiently large, we can obtain a solution of the system of inequalities. Thus, $\mathcal{U}$ is not empty and, by Theorem 4.1, the 2D-RRW-WBP is positive recurrent. Next we prove (C2)-(b). We have $\bar{\mathcal{N}}^{\{1\}} = \{\{1\}, N\}$ and $\bar{\mathcal{N}}_p^{\{1\}} = \{\{1\}, N\}$. Hence, if we find a solution $\boldsymbol{w} = (w_1, w_2)$ to the following system of inequalities, then the solution is an element of $\mathcal{W}_{\{1\}}$ in Theorem 4.2:

$$\langle \boldsymbol{a}(\{1\}), \boldsymbol{w} \rangle = a_1(\{1\})\, w_1 > 0, \quad \langle \boldsymbol{a}(N), \boldsymbol{w} \rangle = a_1(N)\, w_1 + a_2(N)\, w_2 > 0, \quad w_1 > 0,\ w_2 < 0.$$

It is such a solution that $w_1 = 1$ and $w_2 = -1$. Thus, $\mathcal{W}_{\{1\}}$ is not empty and, by Theorem 4.2, the 2D-RRW-WBP is transient. (C3) is symmetric to (C2), thus the proof is analogous.

In (C4), we have $\mathcal{N}_p = \{N\}$ and $\bar{\mathcal{N}}_p^N = \{N\}$. Hence, if we find a solution $\boldsymbol{w} = (w_1, w_2)$ to the following system of inequalities, then the solution is an element of $\mathcal{W}_N$ in Theorem 4.2:

$$\langle \boldsymbol{a}(N), \boldsymbol{w} \rangle = a_1(N)\, w_1 + a_2(N)\, w_2 > 0, \quad w_1 > 0, \quad w_2 > 0.$$

It is such a solution that $w_1 = w_2 = 1$. Thus, $\mathcal{W}_N$ is not empty and, by Theorem 4.2, the 2D-RRW-WBP is transient. □

Remark 5.1. *By Theorem 7.2.3 of Latouche and Rawaswami [7], if* $a_1(N) = 0$ *and* $a_2(N) < 0$*, then* $\mathcal{L}^{\{1\}}$ *is positive recurrent and* $\mathcal{L}^{\{2\}}$ *is not positive recurrent. Thus, we obtain the same result as (C2) in Theorem 4.1. Analogously, if* $a_1(N) < 0$ *and* $a_2(N) = 0$*, then we obtain the same result as (C3) in Theorem 4.1; if (*$a_1(N) > 0$ *and* $a_2(N) = 0$*) or (*$a_1(N) = 0$ *and* $a_2(N) > 0$*) then the 2D-RRW-WBP is transient. However, in the case where* $a_1(N) = 0$ *and* $a_2(N) = 0$*, the 2D-RRW-WBP cannot be classified by our theorems.*

5.2 Three dimensional case

Consider the case of $d = 3$. In this case, we have $N = \{1, 2, 3\}$ and there are seven induced Markov chains: $\mathcal{L}^N$ is a finite Markov chain, $\mathcal{L}^{\{1,2\}}$, $\mathcal{L}^{\{2,3\}}$ and $\mathcal{L}^{\{3,1\}}$ are discrete-time QBD processes, and $\mathcal{L}^{\{1\}}$, $\mathcal{L}^{\{2\}}$ and $\mathcal{L}^{\{3\}}$ are 2D-RRWs-WBP. Hence we can calculate the mean increment vectors $\boldsymbol{a}(N)$, $\boldsymbol{a}(\{1,2\})$, $\boldsymbol{a}(\{2,3\})$ and $\boldsymbol{a}(\{3,1\})$ by standard methods, and classify the induced Markov chains $\mathcal{L}^{\{1\}}$, $\mathcal{L}^{\{2\}}$ and $\mathcal{L}^{\{3\}}$ by Theorem 5.1. However, as far as we know, there are no general methods to obtain the stationary distribution of a 2D-RRW-WBP, and hence it is difficult to obtain the mean increment vectors $\boldsymbol{a}(\{1\})$, $\boldsymbol{a}(\{2\})$ and $\boldsymbol{a}(\{3\})$ when the corresponding induced Markov chains are positive recurrent. Therefore, in the rest of the subsection, supposing that the signs of $\boldsymbol{a}(\{1\})$, $\boldsymbol{a}(\{2\})$ and $\boldsymbol{a}(\{3\})$ are known, we classify 3D-RRWs-WBP.

The classification can be done in a manner similar to that used for the 2D-RRW-WBP, but in this case there are many combinations of the signs of the mean increments. Hence, we summarize only typical cases on Tables 1 to 4, where, for example, "$\boldsymbol{a}(\{1,2\}) = (+\,-\,0)$" means that $a_1(\{1,2\}) > 0$, $a_2(\{1,2\}) < 0$ and $a_3(\{1,2\}) = 0$; "NA" means that the corresponding induced Markov chain is transient, blank means anything, and "positive" ("transient") means that the 3D-RRW-WBP, denoted by $\mathcal{L}$, is positive recurrent (resp. transient). Of course, the tables does not contain all the cases, but every case is shown on some table or symmetric to someone on some tables; for example, the case where $\boldsymbol{a}(N) = (-\,-\,-)$, $\boldsymbol{a}(\{1,2\}) = (-\,-\,0)$, $\boldsymbol{a}(\{2,3\}) = (0\,+\,-)$, $\boldsymbol{a}(\{3,1\}) = (-\,0\,-)$, $\boldsymbol{a}(\{1\}) = (-\,0\,0)$, $\boldsymbol{a}(\{2\}) = (0\,-\,0)$ and $\boldsymbol{a}(\{3\}) = \mathrm{NA}$ is not shown on any table but it is symmetric to Case C1-2-1 on Table 1. We prove only Cases C1-1-1 and C1-1-2; Other cases except for C1-7-1 are analogously proved.

Proof of C1-1-1 and C1-1-2. First we consider C1-1-1. In this case, we have

$$\mathcal{N}_p = \{\{1\}, \{2\}, \{3\}, \{1,2\}, \{2,3\}, \{3,1\}, N\}.$$

In order for a matrix $U = (\boldsymbol{u}_1, \boldsymbol{u}_2, \boldsymbol{u}_3) = (u_{i,j}, i,j = 1,2,3)$ to be positive definite, it suffices that $u_{11} > 0$, $u_{11}u_{22} - u_{12}^2 > 0$ and $\det U > 0$, where we assume $u_{ij} = u_{ji}$ for any i, j. Hence, if we find a solution to the following system of inequalities, then the solution is an element of $\mathcal{U}$ in Theorem 4.1:

$$\begin{aligned}
&\langle \boldsymbol{a}(\{1\}), \boldsymbol{u}_1 \rangle = a_1(\{1\})\, u_{11} < 0,\ \langle \boldsymbol{a}(\{2\}), \boldsymbol{u}_2 \rangle = a_2(\{2\})\, u_{22} < 0,\\
&\langle \boldsymbol{a}(\{3\}), \boldsymbol{u}_3 \rangle = a_3(\{3\})\, u_{33} < 0,\\
&\langle \boldsymbol{a}(\{1,2\}), \boldsymbol{u}_1 \rangle = a_1(\{1,2\})\, u_{11} + a_2(\{1,2\})\, u_{12} < 0,\\
&\langle \boldsymbol{a}(\{1,2\}), \boldsymbol{u}_2 \rangle = a_1(\{1,2\})\, u_{12} + a_2(\{1,2\})\, u_{22} < 0,\\
&\langle \boldsymbol{a}(\{2,3\}), \boldsymbol{u}_2 \rangle = a_2(\{2,3\})\, u_{22} + a_3(\{2,3\})\, u_{23} < 0,\\
&\langle \boldsymbol{a}(\{2,3\}), \boldsymbol{u}_3 \rangle = a_2(\{2,3\})\, u_{23} + a_3(\{2,3\})\, u_{33} < 0,\\
&\langle \boldsymbol{a}(\{3,1\}), \boldsymbol{u}_3 \rangle = a_3(\{3,1\})\, u_{33} + a_1(\{3,1\})\, u_{13} < 0,\\
&\langle \boldsymbol{a}(\{3,1\}), \boldsymbol{u}_1 \rangle = a_3(\{3,1\})\, u_{13} + a_1(\{3,1\})\, u_{11} < 0,\\
&\langle \boldsymbol{a}(N), \boldsymbol{u}_1 \rangle = a_1(N)\, u_{11} + a_2(N)\, u_{12} + a_3(N)\, u_{13} < 0,\\
&\langle \boldsymbol{a}(N), \boldsymbol{u}_2 \rangle = a_1(N)\, u_{12} + a_2(N)\, u_{22} + a_3(N)\, u_{23} < 0,\\
&\langle \boldsymbol{a}(N), \boldsymbol{u}_3 \rangle = a_1(N)\, u_{13} + a_2(N)\, u_{23} + a_3(N)\, u_{33} < 0,\\
&u_{11} > 0,\ u_{11}u_{22} - u_{12}^2 > 0,\\
&u_{11}u_{22}u_{33} + 2\, u_{12}u_{23}u_{13} - u_{11}u_{23}^2 - u_{22}u_{13}^2 - u_{33}u_{12}^2 > 0.
\end{aligned}$$

For a $\delta > 0$, we set $u_{11} = u_{22} = u_{33} = \delta$ and $u_{12} = u_{23} = u_{13} = 1$. Then, since all the mean increments are negative, the inequalities above except for ones on the bottom two lines holds; The fist inequality on the second line from the bottom obviously holds and, from the remaining inequalities, we obtain

$$\delta^2 - 1 > 0, \quad \delta^3 + 2 - 3\delta > 0.$$

For a sufficiently large δ, these two inequalities hold. Hence, $\mathcal{U}$ is not empty and, by Theorem 4.1, the 3D-RRW-WBP is positive recurrent.

Next we consider C1-1-2. We have $\bar{\mathcal{N}}^{\{1\}} = \bar{\mathcal{N}}_p^{\{1\}} = \{\{1\}, \{1,2\}, \{3,1\}, N\}$. Thus, if we find a solution $\boldsymbol{w} = (w_1, w_2, w_3)$ to the following system of inequalities, then the solution is an element of $\mathcal{W}_{\{1\}}$ in Theorem 4.2:

$$\begin{aligned}
&\langle \boldsymbol{a}(\{1\}), \boldsymbol{w} \rangle = a_1(\{1\})\, w_1 > 0,\\
&\langle \boldsymbol{a}(\{1,2\}), \boldsymbol{w} \rangle = a_1(\{1,2\})\, w_1 + a_2(\{1,2\})\, w_2 > 0,\\
&\langle \boldsymbol{a}(\{3,1\}), \boldsymbol{w} \rangle = a_1(\{3,1\})\, w_1 + a_3(\{3,1\})\, w_3 > 0,\\
&\langle \boldsymbol{a}(N), \boldsymbol{w} \rangle = a_1(N)\, w_1 + a_2(N)\, w_2 + a_3(N)\, w_3 > 0,\\
&w_1 > 0,\ w_2 < 0,\ w_3 < 0.
\end{aligned}$$

For a $\delta > 0$, we set $w_1 = \delta$ and $w_2 = w_3 = -1$, then the inequalities are reduced to

$$0 < \delta < \min\left\{ \frac{a_2(\{1,2\})}{a_1(\{1,2\})}, \frac{a_3(\{3,1\})}{a_1(\{3,1\})}, \frac{a_2(N) + a_3(N)}{a_1(N)} \right\},$$

and δ satisfying this inequality exists. Therefore, $\mathcal{W}_{\{1\}}$ is not empty and, by Theorem 4.2, the 3D-RRW-WBP is transient. □

C1-7-1 on Table 1 is a unique case. In relation to semimartingale reflecting Brownian motions (SRBMs), C1-7-1 corresponds to the case where a SRBM has spiral fluid paths (see, for example, Kharroubi et al. [6] and Bramson et al. [2]). The following theorem provides, for C1-7-1, sufficient conditions on which the 3D-RRW-WBP is positive recurrent and on which it is transient; For 3D-RRWs *without a background process*, a similar result was obtained in a different way in Fayolle et al. [5]. For the case where the 3D-RRW-WBP is positive recurrent, the theorem is proved by Theorem 4.1. On the other hand, for the case where the 3D-RRW-WBP is transient, it cannot directly be proved by Theorem 4.2 and some tricks are needed to prove it; Since the proof is complicate and lengthy, we put it in Appendix A.

Theorem 5.2. *(a) Assume the 3D-RRW-WBP satisfies* $\boldsymbol{a}(N) < \mathbf{0}$ *and*

$$a_1(\{1,2\}) < 0,\ a_2(\{1,2\}) > 0,\ a_2(\{2,3\}) < 0,\ a_3(\{2,3\}) > 0,$$
$$a_3(\{3,1\}) < 0,\ a_1(\{3,1\}) > 0.$$

Then it is positive recurrent if

$$\left|\frac{a_2(\{1,2\})}{a_1(\{1,2\})}\right| \left|\frac{a_3(\{2,3\})}{a_2(\{2,3\})}\right| \left|\frac{a_1(\{3,1\})}{a_3(\{3,1\})}\right| < 1, \tag{5.1}$$

and it is transient if

$$\left|\frac{a_2(\{1,2\})}{a_1(\{1,2\})}\right| \left|\frac{a_3(\{2,3\})}{a_2(\{2,3\})}\right| \left|\frac{a_1(\{3,1\})}{a_3(\{3,1\})}\right| > 1. \tag{5.2}$$

(b) C1-7-1: Assume the 3D-RRW-WBP satisfies $\boldsymbol{a}(N) < \mathbf{0}$ *and*

$$a_1(\{1,2\}) > 0,\ a_2(\{1,2\}) < 0,\ a_2(\{2,3\}) > 0,\ a_3(\{2,3\}) < 0,$$
$$a_3(\{3,1\}) > 0,\ a_1(\{3,1\}) < 0.$$

Then it is positive recurrent if

$$\left|\frac{a_1(\{1,2\})}{a_2(\{1,2\})}\right| \left|\frac{a_2(\{2,3\})}{a_3(\{2,3\})}\right| \left|\frac{a_3(\{3,1\})}{a_1(\{3,1\})}\right| < 1, \tag{5.3}$$

and it is transient if

$$\left|\frac{a_1(\{1,2\})}{a_2(\{1,2\})}\right| \left|\frac{a_2(\{2,3\})}{a_3(\{2,3\})}\right| \left|\frac{a_3(\{3,1\})}{a_1(\{3,1\})}\right| > 1. \tag{5.4}$$

Proof of the case where the 3D-RRW-WBP is positive recurrent. We prove only part (a); part (b) is analogously proved. We have

$$\mathcal{N}_p = \{\{1,2\},\{2,3\},\{3,1\},N\}.$$

Hence, for a symmetric matrix $U = (\boldsymbol{u}_1, \boldsymbol{u}_2, \boldsymbol{u}_3) = (u_{i,j})$, if we find a solution to the following system of inequalities, then the solution is an element of $\mathcal{U}$ in Theorem 4.1:

$$\begin{aligned}
&\langle \boldsymbol{a}(\{1,2\}), \boldsymbol{u}_1\rangle = a_1(\{1,2\})\, u_{11} + a_2(\{1,2\})\, u_{12} < 0,\\
&\langle \boldsymbol{a}(\{1,2\}), \boldsymbol{u}_2\rangle = a_1(\{1,2\})\, u_{12} + a_2(\{1,2\})\, u_{22} < 0,\\
&\langle \boldsymbol{a}(\{2,3\}), \boldsymbol{u}_2\rangle = a_2(\{2,3\})\, u_{22} + a_3(\{2,3\})\, u_{23} < 0,\\
&\langle \boldsymbol{a}(\{2,3\}), \boldsymbol{u}_3\rangle = a_2(\{2,3\})\, u_{23} + a_3(\{2,3\})\, u_{33} < 0,\\
&\langle \boldsymbol{a}(\{3,1\}), \boldsymbol{u}_3\rangle = a_3(\{3,1\})\, u_{33} + a_1(\{3,1\})\, u_{13} < 0,\\
&\langle \boldsymbol{a}(\{3,1\}), \boldsymbol{u}_1\rangle = a_3(\{3,1\})\, u_{13} + a_1(\{3,1\})\, u_{11} < 0,\\
&\langle \boldsymbol{a}(N), \boldsymbol{u}_1\rangle = a_1(N)\, u_{11} + a_2(N)\, u_{12} + a_3(N)\, u_{13} < 0,\\
&\langle \boldsymbol{a}(N), \boldsymbol{u}_2\rangle = a_1(N)\, u_{12} + a_2(N)\, u_{22} + a_3(N)\, u_{23} < 0,\\
&\langle \boldsymbol{a}(N), \boldsymbol{u}_3\rangle = a_1(N)\, u_{13} + a_2(N)\, u_{23} + a_3(N)\, u_{33} < 0,\\
&u_{11} > 0,\ u_{11}u_{22} - u_{12}^2 > 0,\\
&u_{11}u_{22}u_{33} + 2\, u_{12}u_{23}u_{13} - u_{11}u_{23}^2 - u_{22}u_{13}^2 - u_{33}u_{12}^2 > 0.
\end{aligned}$$

Assume that $u_{ij} > 0$ for all i, j and define r_{12}, r_{23} and r_{31} as

$$r_{12} = -\frac{a_2(\{1,2\})}{a_1(\{1,2\})} > 0, \quad r_{23} = -\frac{a_3(\{2,3\})}{a_2(\{2,3\})} > 0, \quad r_{31} = -\frac{a_1(\{3,1\})}{a_3(\{3,1\})} > 0,$$

where, by the assumption, we have $r_{12}r_{23}r_{31} < 1$. Then, the inequalities are reduced to

$$\begin{cases} r_{12}\, u_{22} < u_{12} < r_{12}^{-1}\, u_{11}, \quad r_{23}\, u_{33} < u_{23} < r_{23}^{-1}\, u_{22}, \\ r_{31}\, u_{11} < u_{13} < r_{31}^{-1}\, u_{33}, \end{cases} \tag{5.5}$$

$$u_{11}u_{22} - u_{12}^2 > 0, \tag{5.6}$$

$$u_{11}u_{22}u_{33} + 2\, u_{12}u_{23}u_{13} - u_{11}u_{23}^2 - u_{22}u_{13}^2 - u_{33}u_{12}^2 > 0. \tag{5.7}$$

For a δ satisfying $0 < \delta < u_{11}$, set

$$u_{22} = r_{12}^{-2}(u_{11} - \delta), \quad u_{33} = r_{31}^2(u_{11} + \delta). \tag{5.8}$$

Then, letting δ be sufficiently small, we have

$$\frac{r_{12}\, u_{22}}{r_{12}^{-1}\, u_{11}} = \frac{u_{11} - \delta}{u_{11}} < 1, \quad \frac{r_{23}\, u_{33}}{r_{23}^{-1}\, u_{22}} = (r_{12}r_{23}r_{31})^2\, \frac{u_{11} + \delta}{u_{11} - \delta} < 1, \quad \frac{r_{31}\, u_{11}}{r_{31}^{-1}\, u_{33}} = \frac{u_{11}}{u_{11} + \delta} < 1,$$

and this implies that there exist u_{12}, u_{23} and u_{13} satisfying inequalities (5.5). Let c be defined as $c = u_{11}u_{22}u_{33} > 0$ and, for a sufficiently small $\varepsilon > 0$, set

$$u_{12}^2 = (c - \varepsilon)/u_{33}, \quad u_{23}^2 = (c - \varepsilon)/u_{11}, \quad u_{13}^2 = (c - \varepsilon)/u_{22}, \tag{5.9}$$

then these u_{12}, u_{23} and u_{13} satisfy inequalities (5.5) and (5.6). Substituting them into inequality (5.7), we obtain

$$2\sqrt{\frac{(c - \varepsilon)^3}{c}} - 2c + 3\varepsilon > 0, \tag{5.10}$$

and this holds for any $\varepsilon > 0$ since the left hand side of inequality (5.10) is increasing in $\varepsilon \in (0, \infty)$ and takes the value of zero at $\varepsilon = 0$. Therefore, we see that the system of inequalities (5.5) to (5.7) has a solution. Thus, $\mathcal{U}$ is not empty and, by Theorem 4.1, the 3D-RRW-WBP is positive recurrent. □

Table 1: Classification of 3D-RRWs-WBP: $\boldsymbol{a}(N) = (- - -)$.

Case	$\boldsymbol{a}(\{1,2\})$	$\boldsymbol{a}(\{2,3\})$	$\boldsymbol{a}(\{3,1\})$	$\boldsymbol{a}(\{1\})$	$\boldsymbol{a}(\{2\})$	$\boldsymbol{a}(\{3\})$	$\mathcal{L}$
C1-1-1	$(- - 0)$	$(0 - -)$	$(-0-)$	(-00)	$(0 - 0)$	$(00-)$	positive
C1-1-2	$(- - 0)$	$(0 - -)$	$(-0-)$	$(+00)$			transient
C1-1-3	$(- - 0)$	$(0 - -)$	$(-0-)$		$(0 + 0)$		transient
C1-1-4	$(- - 0)$	$(0 - -)$	$(-0-)$			$(00+)$	transient
C1-2-1	$(+ - 0)$	$(0 - -)$	$(-0-)$	(-00)	NA	$(00-)$	positive
C1-2-2	$(+ - 0)$	$(0 - -)$	$(-0-)$	$(+00)$	NA		transient
C1-2-3	$(+ - 0)$	$(0 - -)$	$(-0-)$		NA	$(00+)$	transient
C1-3-1	$(+ + 0)$			NA	NA		transient
C1-4-1	$(+ - 0)$	$(0 + -)$	$(-0-)$	(-00)	NA	NA	positive
C1-4-2	$(+ - 0)$	$(0 + -)$	$(-0-)$	$(+00)$	NA	NA	transient
C1-5-1	$(+ - 0)$	$(0 - +)$	$(-0-)$	(-00)	NA	$(00-)$	positive
C1-5-2	$(+ - 0)$	$(0 - +)$	$(-0-)$	$(+00)$	NA		transient
C1-5-3	$(+ - 0)$	$(0 - +)$	$(-0-)$		NA	$(00+)$	transient
C1-6-1	$(+ - 0)$	$(0 + -)$	$(+0-)$	(-00)	NA	NA	positive
C1-6-1	$(+ - 0)$	$(0 + -)$	$(+0-)$	$(+00)$	NA	NA	transient
C1-7-1	$(+ - 0)$	$(0 + -)$	$(-0+)$	NA	NA	NA	(The. 5.2)

Table 2: Classification of 3D-RRWs-WBP: $\boldsymbol{a}(N) = (+ - -)$.

Case	$\boldsymbol{a}(\{1,2\})$	$\boldsymbol{a}(\{2,3\})$	$\boldsymbol{a}(\{3,1\})$	$\boldsymbol{a}(\{1\})$	$\boldsymbol{a}(\{2\})$	$\boldsymbol{a}(\{3\})$	$\mathcal{L}$
C2-1-1	$(- - 0)$	NA	$(-0-)$	(-00)	$(0 - 0)$	$(00-)$	positive
C2-1-2	$(- - 0)$	NA	$(-0-)$	$(+00)$			transient
C2-1-3	$(- - 0)$	NA	$(-0-)$		$(0 + 0)$		transient
C2-1-4	$(- - 0)$	NA	$(-0-)$			$(00+)$	transient
C2-2-1	$(+ - 0)$	NA	$(-0-)$	(-00)	NA	$(00-)$	positive
C2-2-2	$(+ - 0)$	NA	$(-0-)$	$(+00)$	NA		transient
C2-2-3	$(+ - 0)$	NA	$(-0-)$		NA	$(00+)$	transient
C2-3-1	$(+ + 0)$	NA		NA	NA		transient
C2-4-1	$(+ - 0)$	NA	$(+0-)$	(-00)	NA	NA	positive
C2-4-2	$(+ - 0)$	NA	$(+0-)$	$(+00)$	NA	NA	transient
C2-5-1	$(+ - 0)$	NA	$(-0+)$	NA	NA	$(00-)$	positive
C2-5-2	$(+ - 0)$	NA	$(-0+)$	NA	NA	$(00+)$	transient

Table 3: Classification of 3D-RRWs-WBP: $\boldsymbol{a}(N) = (+\ +\ -)$.

Case	$\boldsymbol{a}(\{1,2\})$	$\boldsymbol{a}(\{2,3\})$	$\boldsymbol{a}(\{3,1\})$	$\boldsymbol{a}(\{1\})$	$\boldsymbol{a}(\{2\})$	$\boldsymbol{a}(\{3\})$	$\mathcal{L}$
C3-1-1	$(-\ -\ 0)$	NA	NA	$(-\,0\,0)$	$(0\ -\ 0)$	NA	positive
C3-1-2	$(-\ -\ 0)$	NA	NA	$(+\,0\,0)$		NA	transient
C3-1-3	$(-\ -\ 0)$	NA	NA		$(0\ +\ 0)$	NA	transient
C3-2-1	$(+\ -\ 0)$	NA	NA	$(-\,0\,0)$	NA	NA	positive
C3-2-2	$(+\ -\ 0)$	NA	NA	$(+\,0\,0)$	NA	NA	transient
C3-3-1	$(+\ +\ 0)$	NA	NA	NA	NA	NA	transient

Table 4: Classification of 3D-RRWs-WBP: $\boldsymbol{a}(N) = (+\ +\ +)$.

Case	$\boldsymbol{a}(\{1,2\})$	$\boldsymbol{a}(\{2,3\})$	$\boldsymbol{a}(\{3,1\})$	$\boldsymbol{a}(\{1\})$	$\boldsymbol{a}(\{2\})$	$\boldsymbol{a}(\{3\})$	$\mathcal{L}$
C4-1-1	NA	NA	NA	NA	NA	NA	transient

5.3 An example of 3D-RRW-WBP

In order to understand a feature of the 3D-RRW-WBP, we consider a simple model of three queues with service interactions (see Fig. 2). Each queue is modeled as an M/M/1 queue with sever vacations and input rejections; When a queue becomes empty, the server of the queue takes an exponentially distributed vacation with the common mean of $1/\delta$ and customers arriving at the queue during the vacation are rejected. After ending the vacation, the sever enters an idle state and waits for customers to arrive. We denote the queues by Q_1, Q_2 and Q_3, respectively, and denote by λ the common arrival rate and by μ the common service rate; We assume $\mu > \lambda$. When all the severs are idle or busy for service, the behavior of the queues are mutually independent. However, when some server takes a vacation, an interaction between the queues occurs as follows: For $l \in \{1,2,3\}$, when the server of Q_l takes a vacation, the server of $\mathrm{Q}_{(l+1) \bmod 3}$ begins to suspend its service; When the server of Q_l ends the vacation and becomes idle, the server of $\mathrm{Q}_{(l+1) \bmod 3}$ resumes its service. Here we consider positive recurrence of this model when the value of δ varies.

Let N be given as $N = \{1,2,3\}$. For $l \in N$, let $X_l(t)$ be the number of customers at Q_l at time t, and define $\boldsymbol{X}(t)$ as $\boldsymbol{X}(t) = (X_l(t),\, l \in N)$. For $l \in N$, let $J_l(t)$ be the state of the server of Q_l at time t, and define $\boldsymbol{J}(t)$ as $\boldsymbol{J}(t) = (J_l(t),\, l \in N)$; "$J_l(t) = 2$" means that the server is serving a customer, "$J_l(t) = 1$" the server is suspending a service, "$J_l(t) = 0$" the server is idle and "$J_l(t) = -1$" the server is on a vacation. Then, $\{\boldsymbol{Y}(t)\} = \{(\boldsymbol{X}(t), \boldsymbol{J}(t))\}$ becomes a continuous-time version of 3D-RRW-WBP, where the state subsets of the background process on the boundary faces except for $\mathcal{B}^{\emptyset}$ are given by

$$
\begin{aligned}
&S^N = \{(2,2,2)\},\ S^{\{1,2\}} = \{(1,2,-1),\ (2,2,0)\},\\
&S^{\{2,3\}} = \{(-1,1,2),\ (0,2,2)\},\ S^{\{3,1\}} = \{(2,-1,1),\ (2,0,2)\},\\
&S^{\{1\}} = \{(1,-1,-1),\ (2,-1,0),\ (1,0,-1),\ (2,0,0)\},\\
&S^{\{2\}} = \{(-1,1,-1),\ (-1,1,0),\ (0,2,-1),\ (0,2,0)\},\\
&S^{\{3\}} = \{(-1,-1,1),\ (-1,0,2),\ (0,-1,1),\ (0,0,2)\}.
\end{aligned}
$$

Let $\mathcal{S}$ be the state space of $\{\boldsymbol{Y}(t)\}$ and, for $\boldsymbol{y}_1, \boldsymbol{y}_2 \in \mathcal{S}$, $\boldsymbol{y}_1 \neq \boldsymbol{y}_2$, let $q(\boldsymbol{y}_1, \boldsymbol{y}_2)$ be the transition rate from $\boldsymbol{y}_1$ to $\boldsymbol{y}_2$. Applying the uniformization technique to $\{\boldsymbol{Y}(t)\}$, we obtain a discrete-time 3D-RRW-WBP, denoted by $\{\boldsymbol{Y}_n\} = \{(\boldsymbol{X}_n, \boldsymbol{J}_n)\}$, where the transition probabilities of $\{\boldsymbol{Y}_n\}$ are given by, for a sufficiently large $\nu > 0$ and for $\boldsymbol{y}_1, \boldsymbol{y}_2 \in \mathcal{S}$,

$$P(\boldsymbol{Y}_{n+1} = \boldsymbol{y}_2 \,|\, \boldsymbol{Y}_n = \boldsymbol{y}_1) = \begin{cases} q(\boldsymbol{y}_1, \boldsymbol{y}_2)/\nu, & \boldsymbol{y}_1 \neq \boldsymbol{y}_2, \\ 1 - \sum_{\boldsymbol{y}' \in \mathcal{S},\, \boldsymbol{y}' \neq \boldsymbol{y}_1} q(\boldsymbol{y}_1, \boldsymbol{y}')/\nu & \boldsymbol{y}_1 = \boldsymbol{y}_2. \end{cases}$$

In this model, the state space of the induced Markov chain $\mathcal{L}^N$ includes only one state, and the mean increments with respect to $\mathcal{L}^N$ are given by $a_1(N) = a_2(N) = a_3(N) = -(\mu - \lambda)/\nu < 0$. Thus, the induced Markov chains $\mathcal{L}^{\{1,2\}}$, $\mathcal{L}^{\{2,3\}}$ and $\mathcal{L}^{\{3,1\}}$ are positive recurrent and they have the same stationary distribution. For example, the stationary distribution of $\mathcal{L}^{\{1,2\}} = \{(\hat{X}_{3,n}, \hat{J}_{3,n})\}$, $\boldsymbol{\pi}^{\{1,2\}} = (p_k,\, k = -1, 0, 1, ...)$, is given by

$$p_{-1} = \frac{1-\rho}{1-\rho+\delta/\lambda}, \quad p_k = \rho^k (\delta/\lambda)\, p_{-1},\ k \geq 0,$$

where $p_{-1} = P(\hat{X}_3 = 0, \hat{J}_3 = -1)$, $p_0 = P(\hat{X}_3 = 0, \hat{J}_3 = 0)$, $p_k = P(\hat{X}_3 = k, \hat{J}_3 = 2)$, $k \geq 1$, and $\rho = \lambda/\mu$; in deriving this stationary distribution, we use the fact that Q_3 is independent of Q_1 and Q_2 in the situation where the induced Markov chain $\mathcal{L}^{\{1,2\}}$ is considered. From the stationary distribution, we obtain the following mean increments:

$$\begin{aligned} &a_1(\{1,2\}) = a_2(\{2,3\}) = a_3(\{3,1\}) = -\big((1-p_{-1})\mu - \lambda\big)/\nu, \\ &a_2(\{1,2\}) = a_3(\{2,3\}) = a_1(\{3,1\}) = -(\mu - \lambda)/\nu < 0, \\ &a_3(\{1,2\}) = a_1(\{2,3\}) = a_2(\{3,1\}) = 0. \end{aligned}$$

First we assume $\delta > \lambda\rho$. Then we have $a_1(\{1,2\}) = a_2(\{2,3\}) = a_3(\{3,1\}) < 0$, and the induced Markov chains $\mathcal{L}^{\{1\}}$, $\mathcal{L}^{\{2\}}$ and $\mathcal{L}^{\{3\}}$ are positive recurrent. The state of the induced Markov chain $\mathcal{L}^{\{1\}}$ consists of the number of customers in Q_2, that in Q_3 and the states of the servers in Q_2 and Q_3. Since service progress in Q_3 may be suspended due to the vacation of the server in Q_2, the stationary probability that the sever is on a vacation in Q_3 is less than or equal to that in isolated Q_3, i.e.,

$$P(\text{the sever is on a vacation in } \mathrm{Q}_3) \leq p_{-1}.$$

Thus, by the assumption that $\delta > \lambda\rho$, we obtain

$$\begin{aligned} a_1(\{1\}) &= -\Big(\big(1 - P(\text{the sever in } \mathrm{Q}_3 \text{ is on a vacation})\big)\mu - \lambda\Big)/\nu \\ &\leq -\big((1-p_{-1})\mu - \lambda\big)/\nu < 0, \end{aligned}$$

and this implies $a_1(\{1\}) = a_2(\{2\}) = a_3(\{3\}) < 0$. Therefore, Case C1-1-1 holds and the 3D-RRW-WBP is positive recurrent. Next we assume $\delta < \lambda\rho$. Then we have $a_1(\{1,2\}) = a_2(\{2,3\}) = a_3(\{3,1\}) > 0$ and the induced Markov chains $\mathcal{L}^{\{1\}}$, $\mathcal{L}^{\{2\}}$ and $\mathcal{L}^{\{3\}}$ are transient. This case, therefore, corresponds to C1-7-1. Thus, if $\lambda\rho > \delta > \lambda\,(\rho - 1/2)$, then we have

$$\left|\frac{a_1(\{1,2\})}{a_2(\{1,2\})}\right| \left|\frac{a_2(\{2,3\})}{a_3(\{2,3\})}\right| \left|\frac{a_3(\{3,1\})}{a_1(\{3,1\})}\right| = \left(\frac{\lambda - (1-p_{-1})\mu}{\mu - \lambda}\right)^3 < 1,$$

and, by Theorem 5.2, the 3D-RRW-WBP is positive recurrent; if $\lambda\,(\rho - 1/2) > \delta$, then it is transient.



Figure 2: A three-queue model with service interactions.

6 Conclusions

We have provided sufficient conditions on which a d-dimensional skip-free RRW with a background process is positive recurrent and on which it is transient, and applying the results, we have classified two and three-dimensional RRWs with a background process. However, our results cannot be applied to the case where the model is probably null recurrent; for example, the case where the mean increment vector $\boldsymbol{a}(N)$ evaluated by the stationary distribution of the induced Markov chain $\mathcal{L}^N$ is equal to a zero vector, i.e., $\boldsymbol{a}(N) = \mathbf{0}$. To understand recurrence and transience of the model in such a case is an interesting problem and we let it be a further study.

References

[1] Bramson, M. (2008). Stability of queueing networks. Probability Survey **5** 169–345.

[2] Bramson, M., Dai, J.G. and Harrison J.M. (2010). Positive recurrence of reflecting Brownian motion in three dimensions. The Annals of Applied Probability **20(2)** 753–783.

[3] Dai, J.G. (1995). On positive harris recurrence of multiclass queueing networks: a unified approach via fluid limit models. The Annals of Applied Probability **5(1)** 49–77.

[4] Fayolle, G. (1989). On random walks arising in queueing systems: ergodicity and transience via quadratic forms as Lyapounov functions – Part I. Queueing Systems **5** 167–184.

[5] Fayolle, G., Malyshev, V.A., and Menshikov, M.V. (1995). *Topics in the Constructive Theory of Countable Markov Chains.* Cambridge University Press, Cambridge.

[6] Kharroubi, A.E., Tahar, A.B. and Yaacoubi, A. (2002). On the stability of the linear Skorohod problem in an orthant. Mathematical Methods of Operations Research **56** 243–258.

[7] Latouche, G. and Ramaswami, V. (1999). *Introduction to Matrix Analytic Methods in Stochastic Modeling.* SIAM, Philadelphia.

[8] Malyshev, V.A. (1993). Networks and dynamical systems. Advances in Applied Probability **25** 140–175.

[9] Meyn, S.P. and Down, D. (1994). Stability of generalized Jackson networks. The Annals of Applied Probability **4(1)** 124–148.

[10] Miyazawa, M. (2011). Light tail asymptotics in multidimensional reflecting processes for queueing networks. TOP **19(2)** 233–299.

[11] Neuts, M.F. (1994). *Matrix-Geometric Solutions in Stochastic Models.* Dover Publications, New York.

[12] Ozawa, T. (2004). Analysis of queues with Markovian Service Processes. Stochastic Models **20(4)** 391–413.

[13] Ozawa, T. (2012). Asymptotics for the stationary distribution in a discrete-time two-dimensional quasi-birth-and-death process. Queueing Systems, to appear (DOI: 10.1007/s11134-012-9323-9).

A Proof of Theorem 5.2

The case where the 3D-RRW-WBP is transient. We prove only part (a); part (b) is analogously proved. First we define a test function. Set c_2 and c_3 so that they satisfy

$$0 < c_2 < \frac{a_2(\{1,2\})}{-a_1(\{1,2\})}, \quad 0 < \frac{c_3}{c_2} < \frac{a_3(\{2,3\})}{-a_2(\{2,3\})}, \quad 0 < \frac{1}{c_3} < \frac{a_1(\{3,1\})}{-a_3(\{3,1\})}; \tag{A.1}$$

It is possible because of inequality (5.2). Let c_0 and c_1 be sufficiently large positive numbers and define vectors $\boldsymbol{w}_{\{1,2\}}$, $\boldsymbol{w}_{\{2,3\}}$ and $\boldsymbol{w}_{\{3,1\}}$ as

$$\begin{aligned} &\boldsymbol{w}_{\{1,2\}} = \frac{1}{c_1}\left(1,\ \frac{1}{c_2},\ \frac{1}{c_3} - c_0\right), \quad \boldsymbol{w}_{\{2,3\}} = \frac{1}{c_1}\left(1 - c_0,\ \frac{1}{c_2},\ \frac{1}{c_3}\right), \\ &\boldsymbol{w}_{\{3,1\}} = \frac{1}{c_1}\left(1,\ \frac{1}{c_2} - c_0,\ \frac{1}{c_3}\right). \end{aligned}$$

In terms of these vectors, we define, for $A \in \{\{1,2\},\{2,3\},\{3,1\}\}$, a function f_A as

$$f_A(\boldsymbol{x}) = \langle \boldsymbol{x}, \boldsymbol{w}_A \rangle,\ \boldsymbol{x} = (x_1, x_2, x_3) \in \mathbb{R}^3.$$

From inequalities (A.1), we obtain

$$f_{\{1,2\}}(\boldsymbol{a}(\{1,2\})) = \frac{1}{c_1}\left(a_1(\{1,2\}) + a_2(\{1,2\})\frac{1}{c_2}\right) > 0, \tag{A.2}$$

$$f_{\{2,3\}}(\boldsymbol{a}(\{2,3\})) = \frac{1}{c_1}\left(a_2(\{2,3\})\frac{1}{c_2} + a_3(\{2,3\})\frac{1}{c_3}\right) > 0, \tag{A.3}$$

$$f_{\{3,1\}}(\boldsymbol{a}(\{3,1\})) = \frac{1}{c_1}\left(a_3(\{3,1\})\frac{1}{c_3} + a_1(\{3,1\})\right) > 0. \tag{A.4}$$

Furthermore, since all the elements of $\boldsymbol{a}(N)$ are negative and c_0 is positive and sufficiently large, we have

$$f_{\{1,2\}}(\boldsymbol{a}(N)) = \frac{1}{c_1}\left(a_1(N) + a_2(N)\frac{1}{c_2} + a_3(N)\frac{1}{c_3} - a_3(N)\,c_0\right) > 0, \tag{A.5}$$

$$f_{\{2,3\}}(\boldsymbol{a}(N)) = \frac{1}{c_1}\left(a_1(N) + a_2(N)\frac{1}{c_2} + a_3(N)\frac{1}{c_3} - a_1(N)\,c_0\right) > 0, \tag{A.6}$$

$$f_{\{3,1\}}(\boldsymbol{a}(N)) = \frac{1}{c_1}\left(a_1(N) + a_2(N)\frac{1}{c_2} + a_3(N)\frac{1}{c_3} - a_2(N)\,c_0\right) > 0, \tag{A.7}$$

where $N = \{1,2,3\}$. We define a positive number ε_0 as

$$\varepsilon_0 = \min_{A\in\{\{1,2\},\{2,3\},\{3,1\}\}} \min\{f_A(\boldsymbol{a}(A)),\ f_A(\boldsymbol{a}(N))\} > 0,$$

which we use after. In order to apply Proposition 3.2, we define a test function f as, for $\boldsymbol{y} = (\boldsymbol{x}, i) \in \mathcal{S}$,

$$f(\boldsymbol{y}) = f(\boldsymbol{x}, i) = \max\left\{f_{\{1,2\}}(\boldsymbol{x}),\ f_{\{2,3\}}(\boldsymbol{x}),\ f_{\{3,1\}}(\boldsymbol{x})\right\},$$

and a subset $\mathcal{A}$ of $\mathcal{S}$ as

$$\mathcal{A} = \{\boldsymbol{y} \in \mathcal{S} : f(\boldsymbol{y}) > 1\}.$$

The plane $f_{\{1,2\}}(\boldsymbol{x}) = 1$ intersects the x_1-axis at $(c_1, 0, 0)$, the x_2-axis at $(0, c_1c_2, 0)$ and the x_3-axis at $(0, 0, -\frac{c_1c_3}{c_0c_3-1})$; the plane $f_{\{2,3\}}(\boldsymbol{x}) = 1$ intersects the x_1-axis at $(-\frac{c_1}{c_0-1}, 0, 0)$, the x_2-axis at $(0, c_1c_2, 0)$ and the x_3-axis at $(0, 0, c_1c_3)$; the plane $f_{\{3,1\}}(\boldsymbol{x}) = 1$ intersects the x_1-axis at $(c_1, 0, 0)$, the x_2-axis at $(0, -\frac{c_1c_2}{c_0c_2-1}, 0)$ and the x_3-axis at $(0, 0, c_1c_3)$. Therefore, since we assumed that c_0 and c_1 are positive and sufficiently large, it can be seen from the definition of the partition $\{\mathcal{V}_A : A \subset N\}$ that we have $\mathcal{A} \subset \mathcal{S} \setminus \mathcal{V}_\emptyset$. It is obvious that $\mathcal{A} \neq \emptyset$ and $\mathcal{A}^C \neq \emptyset$.

Next we consider, for each $\boldsymbol{y} \in \mathcal{S}$, which function is assigned to $f(\boldsymbol{y})$. Define a function $\sigma(\boldsymbol{x})$ as, for $\boldsymbol{x} = (x_1, x_2, x_3) \in \mathbb{R}^3$,

$$\sigma(\boldsymbol{x}) = \begin{cases} \{1,2\}, & x_1 \ge x_3,\ x_2 \ge x_3, \\ \{2,3\}, & x_2 \ge x_1,\ x_3 > x_1, \\ \{3,1\}, & x_3 > x_2,\ x_1 > x_2. \end{cases}$$

Since we have

$$f_{\{2,3\}}(\boldsymbol{x}) - f_{\{1,2\}}(\boldsymbol{x}) = \frac{c_0}{c_1}(x_3 - x_1), \quad f_{\{3,1\}}(\boldsymbol{x}) - f_{\{2,3\}}(\boldsymbol{x}) = \frac{c_0}{c_1}(x_1 - x_2),$$
$$f_{\{1,2\}}(\boldsymbol{x}) - f_{\{3,1\}}(\boldsymbol{x}) = \frac{c_0}{c_1}(x_2 - x_3),$$

$f(\boldsymbol{y})$ is represented as, for $\boldsymbol{y} = (\boldsymbol{x}, i) \in \mathcal{S}$,

$$f(\boldsymbol{y}) = f(\boldsymbol{x}, i) = f_{\sigma(\boldsymbol{x})}(\boldsymbol{x}).$$

Let $\boldsymbol{y} = (\boldsymbol{x}, i)$ be an element of $\mathcal{V}_{\{1,2\}}$, where $\boldsymbol{x} = (x_1, x_2, x_3) \in \mathbb{Z}_+^3$. From the definition of $\mathcal{V}_{\{1,2\}}$, we see that $x_1, x_2 \ge K_{\{1,2\}} > K_N$ and $x_3 \le K_N$. Thus, we have $x_1 > x_3$ and $x_2 > x_3$, and this leads to $\sigma(\boldsymbol{x}) = \{1,2\}$. Analogous results also hold for $\mathcal{V}_{\{2,3\}}$ and $\mathcal{V}_{\{3,1\}}$. As a result, we obtain, for $A \in \{\{1,2\},\{2,3\},\{3,1\}\}$,

$$f(\boldsymbol{y}) = f(\boldsymbol{x}, i) = f_A(\boldsymbol{x}) \text{ if } \boldsymbol{y} = (\boldsymbol{x}, i) \in \mathcal{V}_A. \tag{A.8}$$

Now we come to the phase to evaluate the conditional expectation of the difference between $f(\tilde{\boldsymbol{Y}}_{n+1})$ and $f(\tilde{\boldsymbol{Y}}_n)$, where $\{\tilde{\boldsymbol{Y}}_n\} = \{(\tilde{\boldsymbol{X}}_n, \tilde{J}_n)\}$ is the embedded Markov chain defined in Subsection 4.1.3. Let $\phi(\boldsymbol{y})$ be defined as

$$\phi(\boldsymbol{y}) = E(f(\tilde{\boldsymbol{Y}}_{n+1}) - f(\tilde{\boldsymbol{Y}}_n) \,|\, \tilde{\boldsymbol{Y}}_n = \boldsymbol{y}),\ \boldsymbol{y} \in \mathcal{S},$$

which is represented as

$$\phi(\boldsymbol{y}) = E\Big(f_{\sigma(\tilde{\boldsymbol{X}}_1)}(\tilde{\boldsymbol{X}}_1) - f_{\sigma(\tilde{\boldsymbol{X}}_0)}(\tilde{\boldsymbol{X}}_0) \,|\, \tilde{\boldsymbol{Y}}_0 = \boldsymbol{y}\Big).$$

Since $\{\boldsymbol{X}_n\}$ is skip free in all directions, it is obvios that if $|f(\boldsymbol{y}_1) - f(\boldsymbol{y}_0)| > b$ then we have $P(\boldsymbol{Y}_1 = \boldsymbol{y}_1 \,|\, \boldsymbol{Y}_0 = \boldsymbol{y}_0) = 0$, where

$$b = \frac{2}{c_1}\left(1 + \frac{1}{c_2} + \frac{1}{c_3} + c_0\right).$$

Thus, in order to prove the theorem by Proposition 3.2, it suffices to show that there exists a positive number ε such that $\phi(\boldsymbol{y}) \geq \varepsilon$ for all $\boldsymbol{y} \in \mathcal{A}$. To this end, we divide $\mathcal{A}$ into three parts: $\mathcal{A} \cap \mathcal{V}_N$, $\mathcal{A} \cap \big(\mathcal{V}_{\{1,2\}} \cup \mathcal{V}_{\{2,3\}} \cup \mathcal{V}_{\{3,1\}}\big)$ and $\mathcal{A} \cap \big(\mathcal{V}_{\{1\}} \cup \mathcal{V}_{\{2\}} \cup \mathcal{V}_{\{3\}}\big)$.

(a) Assume $\boldsymbol{y} = (\boldsymbol{x}, i) \in \mathcal{A} \cap \mathcal{V}_N$. Since the induced Markov chain $\mathcal{L}^N$ is positive recurrent, we obtain, by Proposition 4.1,

$$\begin{aligned}\phi(\boldsymbol{y}) &\geq E\Big(f_{\sigma(\boldsymbol{x})}\Big(\sum_{n=1}^{T_N}(\boldsymbol{X}_n - \boldsymbol{X}_{n-1})\Big) \,\Big|\, \tilde{\boldsymbol{Y}}_0 = \boldsymbol{y}\Big) \\ &= T_N\, f_{\sigma(\boldsymbol{x})}\big(\boldsymbol{g}(\boldsymbol{y})\big) \approx T_N\, f_{\sigma(\boldsymbol{x})}\big(\boldsymbol{a}(N)\big) \geq T_N\, \varepsilon_0,\end{aligned}$$

where $\boldsymbol{g}(\boldsymbol{y})$ is given in Subsection 4.2 and we use the fact that $f_{\sigma(\tilde{\boldsymbol{X}}_1)}(\tilde{\boldsymbol{X}}_1) \geq f_{\sigma(\tilde{\boldsymbol{X}}_0)}(\tilde{\boldsymbol{X}}_1)$. This implies that, for any small positive number δ and for a sufficiently large T_N, we have

$$\phi(\boldsymbol{y}) \geq T_N(\varepsilon_0 - \delta) > 0. \tag{A.9}$$

(b) For $A \in \{\{1,2\},\{2,3\},\{3,1\}\}$, assume $\boldsymbol{y} = (\boldsymbol{x}, i) \in \mathcal{A} \cap \mathcal{V}_A$. Since the induced Markov chain $\mathcal{L}^A$ is positive recurrent, we obtain, by Proposition 4.1,

$$\phi(\boldsymbol{y}) \geq E\Big(f_A\Big(\sum_{n=1}^{T_A}(\boldsymbol{X}_n - \boldsymbol{X}_{n-1})\Big) \,\Big|\, \tilde{\boldsymbol{Y}}_0 = \boldsymbol{y}\Big) = T_A\, f_A\big(\boldsymbol{g}(\boldsymbol{y})\big) \approx T_A\, f_A\big(\boldsymbol{a}(A)\big) \geq T_A\, \varepsilon_0.$$

This implies that, for any small positive number δ and for a sufficiently large T_A, we have

$$\phi(\boldsymbol{y}) \geq T_A(\varepsilon_0 - \delta) > 0. \tag{A.10}$$

(c) Assume $\boldsymbol{y} = (\boldsymbol{x}, i) \in \mathcal{A} \cap \mathcal{V}_{\{1\}}$. Since the induced Markov chain $\mathcal{L}_{\{1\}}$ is transient, we use the same technique as that used in the proof of Proposition 4.1. Let $\{(\bar{\boldsymbol{X}}_n, \bar{J}_n)\}$ be the expanded

Markov chain $\bar{\mathcal{L}}_{\{1\}}$ and $\{\tilde{\boldsymbol{Y}}^{\dagger}_{n}\} = \{(\tilde{\boldsymbol{X}}^{\dagger}_{n}, \tilde{J}^{\dagger}_{n})\}$ the embedded Markov chain generated from $\bar{\mathcal{L}}_{\{1\}}$; For the construction of $\{\tilde{\boldsymbol{Y}}^{\dagger}_{n}\}$, especially its embedded points $\{\bar{t}_n\}$, see the proof of Proposition 4.1. For the expanded Markov chain, we modify the functions f and σ and denote them by $\bar{f}$ and $\bar{\sigma}$, respectively; $\bar{f}$ and $\bar{\sigma}$ are given by, for $\boldsymbol{y} = (\boldsymbol{x}, i) \in \bar{\mathcal{S}}_{\{1\}}$,

$$\bar{f}(\boldsymbol{y}) = f_{\bar{\sigma}(\boldsymbol{x})}(\boldsymbol{x}), \quad \bar{\sigma}(\boldsymbol{x}) = \begin{cases} \{1,2\}, & \boldsymbol{y} \in \bar{\mathcal{V}}_{\{1,2\}}, \\ \{3,1\}, & \boldsymbol{y} \in \bar{\mathcal{V}}_{\{3,1\}}, \\ \sigma(\boldsymbol{x}), & \text{otherwise}, \end{cases}$$

where $\bar{\mathcal{S}}_{\{1\}}$ is the state space of $\bar{\mathcal{L}}_{\{1\}}$; $\bar{\mathcal{V}}_{\{1,2\}}$ and $\bar{\mathcal{V}}_{\{3,1\}}$ are the subsets of $\bar{\mathcal{S}}_{\{1\}}$ obtained by expanding $\mathcal{V}_{\{1,2\}}$ and $\mathcal{V}_{\{3,1\}}$, respectively. Then, by Remark 4.2, we have

$$\begin{aligned} \phi(\boldsymbol{y}) &= E\big(\bar{f}(\bar{\boldsymbol{Y}}_{T_{\{1\}}}) - \bar{f}(\bar{\boldsymbol{Y}}_0) \,|\, \bar{\boldsymbol{Y}}_0 = \boldsymbol{y})\big) \\ &= \sum_{n=1}^{T_{\{1\}}} E\Big(1(\bar{t}_n \le T_{\{1\}}) \big(\bar{f}(\tilde{\boldsymbol{Y}}^{\dagger}_{n}) - \bar{f}(\tilde{\boldsymbol{Y}}^{\dagger}_{n-1})\big) \\ &\qquad + 1(\bar{t}_{n-1} \le T_{\{1\}} < \bar{t}_n) \big(\bar{f}(\bar{\boldsymbol{Y}}_{T_{\{1\}}}) - \bar{f}(\tilde{\boldsymbol{Y}}^{\dagger}_{n-1})\big) \,\big|\, \tilde{\boldsymbol{Y}}^{\dagger}_{0} = \boldsymbol{y}\Big) \\ &\ge \sum_{n=1}^{T_{\{1\}}} E\Big(1(\bar{t}_n \le T_{\{1\}})\, f_{\bar{\sigma}(\tilde{\boldsymbol{X}}^{\dagger}_{n-1})}(\tilde{\boldsymbol{X}}^{\dagger}_{n} - \tilde{\boldsymbol{X}}^{\dagger}_{n-1}) \\ &\qquad + 1(\bar{t}_{n-1} \le T_{\{1\}} < \bar{t}_n)\, f_{\bar{\sigma}(\tilde{\boldsymbol{X}}^{\dagger}_{n-1})}(\bar{\boldsymbol{X}}_{T_{\{1\}}} - \tilde{\boldsymbol{X}}^{\dagger}_{n-1}) \,\big|\, \tilde{\boldsymbol{Y}}^{\dagger}_{0} = \boldsymbol{y}\Big) \\ &= T_{\{1\}} \bigg(\sum_{B \in \mathcal{N}^{\{1\}}} \phi_1(B) + \phi_2 \bigg), \end{aligned} \tag{A.11}$$

where $\mathcal{N}^{\{1\}} = \{\{1\}, \{1,2\}, \{3,1\}, N\}$ and, for $B \in \mathcal{N}^{\{1\}}$,

$$\begin{aligned} \phi_1(B) &= \frac{1}{T_{\{1\}}} \sum_{n=1}^{T_{\{1\}}} \sum_{\boldsymbol{y}'=(\boldsymbol{x}',i') \in \bar{\mathcal{V}}_B} E\Big(1(\bar{t}_n \le T_{\{1\}})\, f_{\bar{\sigma}(\boldsymbol{x}')}(\tilde{\boldsymbol{X}}^{\dagger}_{n} - \tilde{\boldsymbol{X}}^{\dagger}_{n-1}) \,\big|\, \tilde{\boldsymbol{Y}}^{\dagger}_{n-1} = \boldsymbol{y}'\Big) \\ &\qquad \times P\big(\tilde{\boldsymbol{Y}}^{\dagger}_{n-1} = \boldsymbol{y}' \,|\, \tilde{\boldsymbol{Y}}^{\dagger}_{0} = \boldsymbol{y}\big), \\ \phi_2 &= \frac{1}{T_{\{1\}}} \sum_{n=1}^{T_{\{1\}}} E\Big(1(\bar{t}_{n-1} \le T_{\{1\}} < \bar{t}_n)\, f_{\bar{\sigma}(\tilde{\boldsymbol{X}}^{\dagger}_{n-1})}(\bar{\boldsymbol{X}}_{T_{\{1\}}} - \tilde{\boldsymbol{X}}^{\dagger}_{n-1}) \,\big|\, \tilde{\boldsymbol{Y}}^{\dagger}_{0} = \boldsymbol{y}\Big). \end{aligned}$$

We have

$$\Big| f_{\sigma(\tilde{\boldsymbol{X}}^{\dagger}_{n-1})}(\bar{\boldsymbol{X}}_{T_{\{1\}}} - \tilde{\boldsymbol{X}}^{\dagger}_{n-1}) \Big| \le \frac{\max\{T_{\{1,2\}}, T_{\{3,1\}}, T_N\}}{c_1} \left(1 + \frac{1}{c_2} + \frac{1}{c_3} + c_0\right), \tag{A.12}$$

and this implies that $\phi_2 \approx 0$ for a sufficiently large $T_{\{1\}}$. Furthermore, we have

$$|\phi_1(\{1\})| \le \frac{1}{c_1} \left(1 + \frac{1}{c_2} + \frac{1}{c_3} + c_0\right) \frac{1}{T_{\{1\}}} \sum_{n=1}^{T_{\{1\}}} P\big(\bar{\boldsymbol{Y}}_{n-1} \in \bar{\mathcal{V}}_{\{1\}} \,|\, \bar{\boldsymbol{Y}}_0 = \boldsymbol{y}\big), \tag{A.13}$$

and this implies that $\phi_1(\{1\}) \approx 0$ for a sufficiently large $T_{\{1\}}$ because the induced Markov chain $\mathcal{L}_{\{1\}}$ is transient. For $B \in \mathcal{N}^{\{1\}} \setminus \{\{1\}\}$, we have

$$\begin{aligned}\phi_1(B) &= \frac{1}{T_{\{1\}}} \sum_{n=1}^{T_{\{1\}}} \sum_{\boldsymbol{y}'=(\boldsymbol{x}',i')\in\bar{\mathcal{V}}_B} T_B\, f_{\bar{\sigma}(\boldsymbol{x}')}(\boldsymbol{g}(\boldsymbol{y}'))\, P\big(\tilde{\boldsymbol{Y}}^{\dagger}_{n-1} = \boldsymbol{y}',\, \bar{t}_n \le T_{\{1\}} \,|\, \tilde{\boldsymbol{Y}}^{\dagger}_0 = \boldsymbol{y}\big) \\ &\approx \frac{T_B}{T_{\{1\}}} \sum_{n=1}^{T_{\{1\}}} \sum_{\boldsymbol{y}'=(\boldsymbol{x}',i')\in\bar{\mathcal{V}}_B} f_{\bar{\sigma}(\boldsymbol{x}')}(\boldsymbol{a}(B))\, P\big(\tilde{\boldsymbol{Y}}^{\dagger}_{n-1} = \boldsymbol{y}',\, \bar{t}_n \le T_{\{1\}} \,|\, \tilde{\boldsymbol{Y}}^{\dagger}_0 = \boldsymbol{y}\big) \\ &\ge \varepsilon_0\, q_{\{1\}}(B), \end{aligned} \tag{A.14}$$

where

$$q_{\{1\}}(B) = \frac{T_B}{T_{\{1\}}} \sum_{n=1}^{T_{\{1\}}} P\big(\tilde{\boldsymbol{Y}}^{\dagger}_{n-1} \in \bar{\mathcal{V}}_B,\, \bar{t}_n \le T_{\{1\}} \,|\, \tilde{\boldsymbol{Y}}^{\dagger}_0 = \boldsymbol{y}\big).$$

Therefore, for any small positive number δ and for a sufficiently large $T_{\{1\}}$, we have

$$\phi(\boldsymbol{y}) \ge T_{\{1\}} \Big(\varepsilon_0\big(q_{\{1\}}(\{1,2\}) + q_{\{1\}}(\{3,1\}) + q_{\{1\}}(N)\big) - \delta\Big) > 0, \tag{A.15}$$

where we use the fact that $q_{\{1\}}(\{1,2\}) + q_{\{1\}}(\{3,1\}) + q_{\{1\}}(N) > 0$ (see expression (4.16) in the proof of Proposition 4.1). Analogously we obtain, for $\boldsymbol{y} \in \mathcal{A} \cap \mathcal{V}_{\{2\}}$ and for a sufficiently large $T_{\{2\}}$,

$$\phi(\boldsymbol{y}) \ge T_{\{2\}} \Big(\varepsilon_0\big(q_{\{2\}}(\{1,2\}) + q_{\{2\}}(\{2,3\}) + q_{\{2\}}(N)\big) - \delta\Big) > 0, \tag{A.16}$$

and for $\boldsymbol{y} \in \mathcal{A} \cap \mathcal{V}_{\{3\}}$ and for a sufficiently large $T_{\{3\}}$,

$$\phi(\boldsymbol{y}) \ge T_{\{3\}} \Big(\varepsilon_0\big(q_{\{3\}}(\{2,3\}) + q_{\{3\}}(\{3,2\}) + q_{\{3\}}(N)\big) - \delta\Big) > 0. \tag{A.17}$$

From the results of (a), (b) and (c), it can be seen that there exists a positive number ε such that, for any $\boldsymbol{y} \in \mathcal{A}$, $\phi(\boldsymbol{y}) \ge \varepsilon$, and this completes the proof. □